

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 April 2001 (26.04.2001)

PCT

(10) International Publication Number
WO 01/29261 A2

- (51) International Patent Classification⁷: **C12Q 1/68** **WEI, Lun-Yan** [US/US]; 13 Pine Tree Place, West Caldwell, NJ 07006 (US).
- (21) International Application Number: PCT/US00/28352
- (22) International Filing Date: 13 October 2000 (13.10.2000)
- (74) Agent: **FELTHAM, S., Neil**; E.I. du Pont de Nemours and Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).
- (25) Filing Language: English
- (81) Designated States (*national*): AU, CA, CN, IL, IN, JP, MX, NZ, SG, US.
- (26) Publication Language: English
- (84) Designated States (*regional*): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
- (30) Priority Data:
60/159,898 15 October 1999 (15.10.1999) US
- (71) Applicant (*for all designated States except US*): **E.I. DU PONT DE NEMOURS AND COMPANY** [US/US]; 1007 Market Street, Wilmington, DE 19898 (US).
- Published:
— Without international search report and to be republished upon receipt of that report.
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **LAROSSA, Robert** [US/US]; 672 Kadar Drive, West Chester, PA 19382 (US).
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: A METHOD FOR HIGH-DENSITY MICROARRAY MEDIATED GENE EXPRESSION PROFILING

(57) Abstract: The global effect on genes under different environmental conditions can be determined by a comprehensive gene expression profile. The present invention provides a method to monitor the changes in comprehensive cellular gene expression levels at single length resolution by using a high-density microarray prepared with a comprehensive collection of ORFs of a genome. Under different environmental conditions, directly and indirectly affected genes can be detected as the gene expression levels are induced or repressed in comparison to the control.

WO 01/29261 A2

TITLEA METHOD FOR HIGH-DENSITY MICROARRAY MEDIATED
GENE EXPRESSION PROFILING

This application claims the benefit of U.S. Provisional Application
5 No. 60/159,898, filed October 15, 1999.

FIELD OF THE INVENTION

This invention is in the field of bacterial gene expression. More
specifically, this invention is a method for the high density, microarray-mediated
gene expression profiling of *Escherichia coli* for comprehensive gene expression
10 analysis.

BACKGROUND OF THE INVENTION

Escherichia coli has been exhaustively studied for over 50 years. Early
experiments measured the molecular fluxes from small compounds into
macromolecular constituents. These studies were followed by others in which
15 small molecule pools of central metabolic building blocks, nucleotides and amino
acids were determined. The levels of several macromolecular components,
including individual species of proteins, have been measured. Such measurements
of the steady state provide a census of the cellular content while changes upon
imposition of a stress catalogue the cell's fight for survival. This response to an
20 insulting or adverse condition can take many forms from relieving end product
inhibition to derepressing transcription.

In *E. coli*, experiments to define stress-related, global regulatory responses
have often relied upon one of two approaches. In the first, operon fusions induced
by a particular stress are isolated. In the second, proteomic measures in which the
25 protein fractions from stressed and un-stressed cultures are separated by a
two-dimensional method and then compared. Each method has an inherent
technological hurdle; for the former, the map location of responsive gene fusions
must be known precisely, and for the latter, induced or repressed proteins excised
from the two-dimensional gels must be identified.

30 Another method uses a transposon-mediated mutagenesis (Spector et al. *J.*
Bacteriol. 170:345-351 (1988)). A reporter gene is inserted at a random location
in the genome using a transposon. By assaying for the reporter gene before and
after the treatment, genes affected by the treatment can be mapped and cloned by
using the linked transposon as a marker. However, this method is limited to
35 non-essential genes.

Alternatively, mRNA measurements utilizing techniques (such as
hybridization to DNA and primer extension) have allowed the monitoring of

individual gene's expression profiles. DeRisi et al. (*Science* 278:680-686 (1997)) reported the expression profiling of most yeast genes. The measurements were facilitated by high-density arrays of individual genes and specific labeling of cDNA copies of eukaryotic mRNA using polyA tail-specific primers. The lack of a polyA tail and the extremely short bacterial mRNA half life represent hurdles for the application of DNA micro-array technology to prokaryotic research.

A comprehensive expression profiling has been performed previously with the yeast *Saccharomyces cerevisiae*. Adaptation of RNA isolation and labeling protocols from eukaryotes to prokaryotes is not straightforward since eukaryotic mRNA manipulations often exploit 3'-polyadenylation of this molecular species.

Chuang et al. (*J. Bacteriol.* 175:2026-2036 (1993)) reported an expression profiling using large DNA fragments from an ordered λ library of *E. coli* genomic fragments as a capture reagent. It allowed the comparison of the expression patterns from large portions of DNA fragments by comparing mRNA levels from stressed and unstressed *E. coli* cultures. The resolution of this method, however, was unsatisfactory. Expression of groups of genes, as opposed to the expression of each individual gene was measured. Moreover, the method used radio-labeled DNA as a probe with the incumbent need for safety precautions. Furthermore, the use of radio-labeled probe prevents the simultaneous measurement of the expression level in a test sample and a control sample.

Richmond et al. (*Nucleic Acids Research*, 19:3821-3835 (1999)) has recently reported genome-wide expression profiling of *E. coli* at a single ORF level of resolution. Changes in RNA levels after exposure to heat shock or IPTG were analyzed using comprehensive low density blots of individual ORFs on a nylon matrix and comprehensive high density arrays of individual ORFs spotted on glass slides. The results of the two methods were compared.

The methods recited above permit monitoring of the effect of environmental changes on gene expression by comparing expression levels of a limited number of genes. They, however, fail to monitor the comprehensive responses of a preponderance of individual genes in the genome of an organism in reliable, useful manner.

The problem to be solved, therefore, is to provide a way to measure the comprehensive gene expression profile analysis of the organism.

SUMMARY OF THE INVENTION

The invention provides a method for identifying gene expression changes within a bacterial species comprising:

- (a) providing a comprehensive micro-array synthesized from DNA comprised in a bacterial species;

- 5
- 10
- 15
- 20
- (b) generating a first set of labeled probes from bacterial RNA, the RNA isolated from the bacterial species of step (a);
 - (c) hybridizing the first set of labeled probes of step (b) to the comprehensive micro-array of step (a), wherein hybridization results in a detectable signal generated from the labeled probe;
 - (d) measuring the signal generated by the hybridization of the first set of labeled probe to the comprehensive micro-array of step (c);
 - (e) subjecting the bacterial species of step (a) to a gene expression altering condition whereby the gene expression profile of the bacterial species is altered to produce a modified bacterial species ;
 - (f) generating a second set of labeled probes from bacterial RNA, the RNA isolated from the modified bacterial species of step (e);
 - (g) hybridizing the second set of labeled probes of step (f) to the comprehensive micro-array of step (a), wherein hybridization results in a detectable signal generated from the labeled probe;
 - (h) measuring the signal generated by the hybridization of the second set of labeled probes to the comprehensive micro-array of step (g); and
 - (i) comparing signal generated from the first hybridization to the signal generated from the second hybridization to identify gene expression changes within a bacterial species.

Additionally the invention provides a method for identifying gene expression changes within a bacterial strain comprising:

- 25
- 30
- 35
- (a) providing a comprehensive micro-array synthesized from DNA comprised in a bacterial species ;
 - (b) generating a first set of fluorescent cDNA from bacterial RNA, the RNA isolated from the bacterial species of step (a);
 - (c) hybridizing the first set of fluorescent cDNA of step (b) to the comprehensive micro-array of step (a), wherein hybridization results in a detectable signal generated from the fluorescent cDNA;
 - (d) measuring the signal generated by the hybridization of the first set of fluorescent cDNA to the comprehensive micro-array of step (c);
 - (e) subjecting the bacterial species of step (a) to a gene expression altering condition whereby the gene expression profile of the bacterial species is altered to produce a modified bacterial species;
 - (f) generating a second set of fluorescent cDNA from bacterial RNA, the RNA isolated from the modified bacterial species of step (e);

- (g) hybridizing the second set of fluorescent cDNA of step (f) to the comprehensive micro-array of step (a), wherein hybridization results in a detectable signal generated from the fluorescent cDNA;
- (h) measuring the signal generated by the hybridization of the second set of fluorescent cDNA to the comprehensive micro-array of step (g); and
- (i) comparing signal generated from the first hybridization to the signal generated from the second hybridization to identify gene expression changes within a bacterial species .

10 In an alternate embodiment the invention provides a method for identifying gene expression changes within a genome comprising:

- (a) providing a comprehensive micro-array synthesized from DNA comprised in a prokaryotic or eukaryotic species;
- (b) generating a control set of fluorescent cDNA from total or polyadenylated RNA, the RNA isolated from the species of step (a), the fluorescent cDNA comprising at least one first fluorescent label and at least one different second fluorescent label;
- (c) mixing the control set of fluorescent cDNA labeled with the at least one first label with the control set of fluorescent cDNA labeled with the at least second first label to for a dual labeled control cDNA;
- (d) hybridizing the dual labeled control set of fluorescent cDNA of step (c) to the comprehensive micro-array of step (a), wherein hybridization results in a detectable signal generated from the fluorescent cDNA;
- (e) measuring the signal generated by the hybridization of the dual labeled control set of fluorescent cDNA to the comprehensive micro-array of step (c);
- (f) subjecting the prokaryote or eukaryote of step (a) to a gene expression altering condition whereby the gene expression profile of the prokaryote or eukaryote is altered to produce a modified prokaryote or eukaryote ;
- (g) generating an experimental set of fluorescent cDNA from total or polyadenylated RNA, the RNA isolated from the modified prokaryote or eukaryote of step (e), the fluorescent cDNA comprising the first fluorescent label and the different second fluorescent label to step (b);

- 5 (h) mixing the experimental set of fluorescent cDNA labeled with the at least one first label with the experimental set of fluorescent cDNA labeled with the at least second first label to form a dual labeled experimental cDNA;
- (i) hybridizing the experimental set of fluorescent cDNA of step (h) to the comprehensive micro-array of step (a), wherein hybridization results in a detectable signal generated from the fluorescent cDNA;
- 10 (j) measuring the signal generated by the hybridization of the second set of fluorescent cDNA to the comprehensive micro-array of step (g); and
- (k) comparing signal generated from the dual labeled control hybridization with the dual labeled experimental hybridization to identify gene expression changes within a prokaryotic or eukaryotic species.
- 15 In another embodiment the invention provides a method for quantitating the amount of protein specifying RNA contained within a genome comprising:
- (a) providing a comprehensive micro-array comprising a multiplicity of genes synthesized from genomic DNA comprised in a prokaryotic or eukaryotic organism;
- 20 (b) generating a set of fluorescent cDNA from total or poly-adenylated RNA isolated from the prokaryotic or eukaryotic organism of step (a);
- (c) generating a set of fluorescent DNA from genomic DNA isolated from the prokaryotic or eukaryotic organism of step (a);
- 25 (d) hybridizing the fluorescent cDNA of step (b) to the comprehensive micro-array of step (a), wherein hybridization results in a first fluorescent signal generated from the fluorescent cDNA for each gene;
- (e) hybridizing the fluorescent DNA of step (c) to the comprehensive micro-array of step (a), wherein hybridization results in a second fluorescent signal generated from the fluorescent DNA for each gene; and
- 30 (f) dividing, for each open reading from, the first fluorescent signal into the second fluorescent signal to provide a quantitated measure of the amount of protein specifying RNA for each gene.
- 35

The methods of the present invention are applicable to genomes contained within a variety of organisms including bacteria, cyanobacteria, yeasts, filamentous fungi, plant cells and animal cells.

The present methods of identifying gene expression changes within genome may be additionally coupled with the methods of quantitating the amount of protein specifying RNA contained within a genome as disclosed herein.

BRIEF DESCRIPTION OF THE DRAWINGS AND

5

SEQUENCE DESCRIPTIONS.

Figure 1A describes the gene expression analysis of IPTG induction in a single hybridization experiment using different slide sets as capture reagents for Cy3-labeled cDNA derived from treated and control cells and plotted in log-log form.

10

Figure 1B describes the gene expression analysis of IPTG induction by labeling the control sample with Cy5 and the induced sample with Cy3 before hybridizing to a single set of 3 slides.

Figure 1C describes an average of induced RNA and control RNA with Cy3 from IPTG induction, generated by label swapping.

15

Figure 1D describes data replicating the results shown in Figure 1C.

Figure 1E describes an averaging of the data of Figure 1C and Figure 1D.

Figure 2 describes the distribution of gene expression levels for cells grown in minimal or rich medium.

20

Figure 3 describes the fractional (summed open reading frame transcripts/total open reading frame transcripts) analysis of gene expression.

The invention can be more fully understood from the following detailed description and the accompanying sequence descriptions which form a part of this application.

25

The following sequences comply with 37 C.F.R. 1.821-1.825 ("Requirements for Patent Applications Containing Nucleotide Sequences and/or Amino Acid Sequence Disclosures - the Sequence Rules") and are consistent with World Intellectual Property Organization (WIPO) Standard ST.25 (1998) and the sequence listing requirements of the EPO and PCT (Rules 5.2 and 49.5(a-bis), and Section 208 and Annex C of the Administrative Instructions). The symbols and format used for nucleotide and amino acid sequence data comply with the rules set forth in 37 C.F.R. §1.822.

30

SEQ ID NO:1 and 2 are primers used in the amplification of the *sdiA* gene.

DETAILED DESCRIPTION OF THE INVENTION

35

Applicants have solved the stated problem by providing a method to measure a comprehensive mRNA expression of *E. coli* using a high density DNA microarray with a near-complete collection of *E. coli* open reading frames (ORFs).

The present invention advances the art by providing:

(i) the first instance of a comprehensive micro-array comprising greater than 75% of all open reading frames from a prokaryotic organism, overcoming the problems of high concentration of endogenous RNAase and ribosomal RNA;

5 (ii) a method for quantitating the amount of each protein specifying RNA contained within a culture; and

(iii) a method for decreasing the background noise generated within a gene expression profile through the combination of multiple signal generating labels.

The present invention has utility in many different fields. Many discovery compounds can be screened by comparing their gene expression profile to a
10 known compound that affects the desirable target gene products. Additionally gene expression profiles are good indicators of genotypic alterations among strains. The present invention may allow the discovery of complementary target inhibitors in combination drug-therapy and may be used as a modeling system to test perturbations in process conditions to determine the conditions for the high
15 yield of desired production in various bio-processes and biotransformations.

In this disclosure, a number of terms and abbreviations are used. The following definitions are provided.

"Open reading frame" is abbreviated ORF. The term "ORF" is refers to a gene that specifies a protein.

20 "Polymerase chain reaction" is abbreviated PCR.

The term "micro-array" means an array of regions having a density of discrete regions of oligonucleotides of at least about 100/cm², and preferably at least about 1000/cm².

25 The term "comprehensive micro array" refers to high-density micro-array containing at least 75% of all open reading frames of the organism.

The term "expression profile" refers to the expression of groups of genes.

The term "gene expression profile" refers to the expression of an individual gene and of suites of individual genes.

30 The "comprehensive expression profile" refers to the gene expression profile of more than 75% of all genes in the genome.

The term "high density" as used in conjunction with micro-array means and array having an array density of generally greater than about 60, more generally greater than about 100, most generally greater than about 600, often greater than about 1000, more often greater than about 5,000, most often greater
35 than about 10,000, preferably greater than about 40,000 more preferably greater than about 100,000, and most preferably greater than about 400,000 different nucleic acids per cm.²

As used herein, an "isolated nucleic acid fragment" is a polymer of RNA or DNA that is single- or double-stranded, optionally containing synthetic, non-natural or altered nucleotide bases. An isolated nucleic acid fragment in the form of a polymer of DNA may be comprised of one or more segments of cDNA,
5 genomic DNA or synthetic DNA.

The term "probe" refers to a single-stranded nucleic acid molecule that can base pair with a complementary single stranded target nucleic acid to form a double-stranded molecule.

The term "genotype" refers to the genetic constitution of an organism as
10 distinguished from its physical appearance.

The term "genomic DNA" refers to total DNA from an organism.

The term "total RNA" refers to non-fractionated RNA from an organism.

The term "protein specifying RNA" or "protein specifying transcript" or "mRNA" refers to RNA derived from ORF.

15 The term "label" will refer to a substance which may be incorporated into DNA or RNA which will emit a detectable signal under various conditions. Typically a label will be a fluorescent moiety.

A nucleic acid molecule is "hybridizable" to another nucleic acid molecule, such as a cDNA, genomic DNA, or RNA, when a single stranded form
20 of the nucleic acid molecule can anneal to the other nucleic acid molecule under the appropriate conditions of temperature and solution ionic strength. Hybridization and washing conditions are well known and exemplified in Sambrook, J., Fritsch, E. F. and Maniatis, T. Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring
25 Harbor (1989), particularly Chapter 11 and Table 11.1 therein. The conditions of temperature and ionic strength determine the "stringency" of the hybridization. Hybridization requires that the two nucleic acids contain complementary sequences, although depending on the stringency of the hybridization, mismatches between bases are possible. The appropriate stringency for hybridizing nucleic
30 acids depends on the length of the nucleic acids and the degree of complementation, variables well known in the art. The greater the degree of similarity or homology between two nucleotide sequences, the greater the value of T_m for hybrids of nucleic acids having those sequences. The relative stability (corresponding to higher T_m) of nucleic acid hybridizations decreases in the
35 following order: RNA:RNA, DNA:RNA, DNA:DNA. For hybrids of greater than 100 nucleotides in length, equations for calculating T_m have been derived (see Sambrook et al., *supra*, 9.50-9.51). For hybridizations with shorter nucleic acids, i.e., oligonucleotides, the position of mismatches becomes more important,

and the length of the oligonucleotide determines its specificity (see Sambrook et al., *supra*, 11.7-11.8). Furthermore, the skilled artisan will recognize that the temperature and wash solution salt concentration may be adjusted as necessary according to factors such as length of the probe.

- 5 The term "complementary" is used to describe the relationship between nucleotide bases that are capable to hybridizing to one another. For example, with respect to DNA, adenosine is complementary to thymine and cytosine is complementary to guanine.

- 10 "Gene" refers to the part of the genome specifying a macromolecular product be it RNA or a protein and include regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence.

A "genetic site" refers to a genomic region at which a gene product operates.

- 15 "Coding sequence" or "open reading frame" (ORF) refers to a DNA sequence that codes for a specific amino acid sequence. "Suitable regulatory sequences" refer to nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences may include
20 promoters, translation leader sequences, introns, and polyadenylation recognition sequences.

- "Promoter" refers to a DNA sequence capable of controlling the expression of a coding sequence or functional RNA. In general, a coding
25 sequence is located 3' to a promoter sequence. Promoters may be derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, or even comprise synthetic DNA segments. It is understood by those skilled in the art that different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of
30 development, or in response to different environmental conditions. Promoters which cause a gene to be expressed in most cell types at most times are commonly referred to as "constitutive promoters". It is further recognized that since in most cases the exact boundaries of regulatory sequences have not been completely defined, DNA fragments of different lengths may have identical promoter activity.

- 35 "RNA transcript" refers to the product resulting from RNA polymerase-catalyzed transcription of a DNA sequence. When the RNA transcript is the polymer product of an RNA polymerase, it is referred to as the primary transcript or it may be a RNA sequence derived from post-transcriptional processing of the

primary transcript and is referred to as the mature RNA. "Messenger RNA (mRNA)" refers to the RNA that is without introns and that can be translated into protein by the cell. "cDNA" refers to a double-stranded DNA that is complementary to and derived from mRNA.

- 5 The term "expression", as used herein, refers to the transcription and stable accumulation of sense (mRNA) or antisense RNA derived from genomic DNA. Expression may also refer to translation of mRNA into a polypeptide.

 The term "stress" or "environmental stress" refers to the condition produced in a cell as the result of exposure to an environmental insult.

- 10 The term "insult" or "environmental insult" refers to any substance or environmental change that results in an alteration of normal cellular metabolism in a bacterial cell or population of cells. Environmental insults may include, but are not limited to, chemicals, environmental pollutants, heavy metals, changes in temperature, changes in pH, as well as agents producing oxidative damage, DNA
- 15 damage, anaerobiosis, and changes in nitrate availability or pathogenesis.

 The term "stress response" refers to the cellular response to an environmental insult.

 The term "stress gene" refers to any gene whose transcription is induced as a result of environmental stress or by the presence of an environmental insult.

- 20 The term "modified bacterial species" refers to a bacterial culture that has been exposed to a stress or insult such that either it demonstrates a change in its gene expression profile. Typically the modified bacterial species is produced as the result of induction or challenge of the culture with a chemical or environmental challenge. Similarly, a "modified prokaryotic or eukaryotic
- 25 species" refers to either a prokaryotic or eukaryotic organism that has been exposed to a stress or insult such that the gene expression profile of that organisms as been altered.

- The term "log phase", "log phase growth", "exponential phase" or "exponential phase growth" refers to cell cultures of organisms growing under
- 30 conditions permitting the exponential multiplication of the cell number.

- The term "growth-altering environment" refers to energy, chemicals, or living things that have the capacity to either inhibit cell growth or kill cells. Inhibitory agents may include but are not limited to mutagens, antibiotics, UV light, gamma-rays, x-rays, extreme temperature, phage, macrophages, organic
- 35 chemicals and inorganic chemicals.

 Standard recombinant DNA and molecular cloning techniques used here are well known in the art and are described by Sambrook, J., Fritsch, E. F. and Maniatis, T., Molecular Cloning: A Laboratory Manual, Second Edition, Cold

Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1989) (hereinafter "Maniatis"); and by Silhavy, T. J., Bennis, M. L. and Enquist, L. W., Experiments with Gene Fusions, Cold Spring Harbor Laboratory Cold Press Spring Harbor, NY (1984); and by Ausubel, F. M. et al., Current Protocols in Molecular Biology, published by Greene Publishing Assoc. and Wiley-Interscience (1987).

The present invention provides a method to measure the changes in gene expression profiles of prokaryotic organisms. The present invention also provides a method to measure the levels of protein specifying RNA in prokaryotic and/or eukaryotic organisms. The present invention provides a method to compare the gene expression patterns of two samples differing in one variable. The variables may include but are not limited to genotype, media, temperature, depletion or addition of nutrient, addition of an inhibitor, physical assault, irradiation, heat, cold, elevated or lowered pressure, desiccation, low or high ionic strength, and growth phases.

Gene expression profiles were determined under the following conditions to find: (a) differences in gene expression profiles caused by growth of *E. coli* in either minimal or rich medium, (b) changes in gene expression associated with the transition from exponential phase to stationary phase growth in minimal medium, and (c) the specificity of induction mediated by isopropylthiogalactoside (IPTG), the classic *lac* operon inducer, (d) the specificity of expression changes mediated by the amplification of *sdiA*, a positive activator of an operon that includes *ftsQAZ*, genes essential for septation, and (e) the changes in gene expression patterns with cells that cannot turn on the SOS stress response in comparison to wild type response when the cells are exposed to mitomycin C (MMC).

In its most basic form the present invention creates a comprehensive micro-array from a bacterial genome. Any bacteria is suitable for analysis by the method of the present invention where enteric bacteria (*Escherichia*, and *Salmonella* for example) as well as cyanobacteria (such as *Rhodobacter* and *Synechocystis* and *Bacillus*, *Acinetobacter*, *Streptomyces*, *Methylobacter*, and *Pseudomona* are particularly suitable.

One of skill in the art will appreciate that in order to measure the transcription level (and thereby the expression level) of a gene or genes, it is desirable to provide a nucleic acid sample comprising mRNA transcript(s) of the gene or genes, or nucleic acids derived from the mRNA transcript(s). As used herein, a nucleic acid derived from an mRNA transcript refers to a nucleic acid for whose synthesis the mRNA transcript or a subsequence thereof has ultimately served as a template. Thus, a cDNA reverse transcribed from an mRNA, an RNA

transcribed from that cDNA, a DNA amplified from the cDNA, an RNA transcribed from the amplified DNA, etc., are all derived from the mRNA transcript and detection of such derived products is indicative of the presence and/or abundance of the original transcript in a sample. Thus, suitable samples
5 include, but are not limited to, mRNA transcripts of the gene or genes, cDNA reverse transcribed from the mRNA, cRNA transcribed from the cDNA, DNA amplified from the genes, RNA transcribed from amplified DNA, and the like.

Typically the genes are amplified by methods of primer directed amplification such as polymerase chain reaction (PCR) (U.S. Patent
10 No. 4,683,202 (1987, Mullis, et al.) and U.S. Patent No. 4,683,195 (1986, Mullis, et al.), ligase chain reaction (LCR) (Tabor et al., *Proc. Acad. Sci. U.S.A.*, 82, 1074-1078 (1985)) or strand displacement amplification (Walker et al., *Proc. Natl. Acad. Sci. U.S.A.*, 89, 392, (1992) for example.

The micro-array is comprehensive in that it incorporates at least 75% of all
15 ORF's present in the genome. Amplified ORF's are then spotted on slides comprised of glass or some other solid substrate by methods well known in the art to form a micro-array. Methods of forming high density arrays of oligonucleotides, with a minimal number of synthetic steps are known (see for example Brown et al., U.S. Patent No. 6,110,426). The oligonucleotide analogue
20 array can be synthesized on a solid substrate by a variety of methods, including, but not limited to, light-directed chemical coupling, and mechanically directed coupling. See Pirrung et al., U.S. Pat. No. 5,143,854 (see also PCT Application No. WO 90/15070) and Fodor et al., PCT Publication Nos. WO 92/10092 and WO 93/09668 which disclose methods of forming vast arrays of peptides,
25 oligonucleotides and other molecules using, for example, light-directed synthesis techniques. See also, Fodor et al., *Science*, 251, 767-77 (1991).

Bacteria typically contain from about 2000 to about 6000 ORF's per genome and the present method is suitable for genomes of this size where genomes of about 4000 ORF's are most suitable. The ORF's are arrayed in high
30 density on at least one glass microscope slide. This is in contrast to a low density array where ORF's are arrayed on a membranous material such as nitrocellulose. The small surface area of the high density array (often less than about 10 cm², preferably less than about 5 cm² more preferably less than about 2 cm², and most preferably less than about 1.6 cm²) permits extremely uniform hybridization
35 conditions (temperature regulation, salt content, etc.).

Once all the genes of ORF's from the genome are amplified, isolated and arrayed, a set of probes, bearing a signal generating label are synthesized. Probes may be randomly generated or may be synthesized based on the sequence of

specific open reading frames. Probes of the present invention are typically single stranded nucleic acid sequences which are complementary to the nucleic acid sequences to be detected. Probes are "hybridizable" to the ORF's. The probe length can vary from 5 bases to tens of thousands of bases, and will depend upon the specific test to be done. Typically a probe length of about 15 bases to about 30 bases is suitable. Only part of the probe molecule need be complementary to the nucleic acid sequence to be detected. In addition, the complementarity between the probe and the target sequence need not be perfect. Hybridization does occur between imperfectly complementary molecules with the result that a certain fraction of the bases in the hybridized region are not paired with the proper complementary base.

Signal generating labels that may be incorporated into the probes are well known in the art. For example labels may include but are not limited to fluorescent moieties, chemiluminescent moieties, particles, enzymes, radioactive tags, or light emitting moieties or molecules, where fluorescent moieties are preferred. Most preferred are fluorescent dyes capable of attaching to nucleic acids and emitting a fluorescent signal. A variety of dyes are known in the art such as fluorescein, Texas red, and rhodamine. Preferred in the present invention are the mono reactive dyes cy3 (146368-16-3) and cy5 (146368-14-1) both available commercially (i.e. Amersham Pharmacia Biotech, Arlington Heights, IL). Suitable dyes are discussed in U.S. Patent No. 5,814,454 hereby incorporated by reference.

Labels may be incorporated by any of a number of means well known to those of skill in the art. However, in a preferred embodiment, the label is simultaneously incorporated during the amplification step in the preparation of the probe nucleic acids. Thus, for example, polymerase chain reaction (PCR) with labeled primers or labeled nucleotides will provide a labeled amplification product. In a preferred embodiment, reverse transcription or replication, using a labeled nucleotide (e.g. dye-labeled UTP and/or CTP) incorporates a label into the transcribed nucleic acids.

Alternatively, a label may be added directly to the original nucleic acid sample (e.g., mRNA, polyA mRNA, cDNA, etc.) or to the amplification product after the synthesis is completed. Means of attaching labels to nucleic acids are well known to those of skill in the art and include, for example nick translation or end-labeling (e.g. with a labeled RNA) by kinasing of the nucleic acid and subsequent attachment (ligation) of a nucleic acid linker joining the sample nucleic acid to a label (e.g., a fluorophore).

Following incorporation of the label into the probe the probes are then hybridized to the micro-array using standard conditions where hybridization results in a double stranded nucleic acid, generating a detectable signal from the label at the site of capture reagent attachment to the surface. Typically the probe and array must be mixed with each other under conditions which will permit nucleic acid hybridization. This involves contacting the probe and array in the presence of an inorganic or organic salt under the proper concentration and temperature conditions. The probe and array nucleic acids must be in contact for a long enough time that any possible hybridization between the probe and sample nucleic acid may occur. The concentration of probe or array in the mixture will determine the time necessary for hybridization to occur. The higher the probe or array concentration the shorter the hybridization incubation time needed. Optionally a chaotropic agent may be added. The chaotropic agent stabilizes nucleic acids by inhibiting nuclease activity. Furthermore, the chaotropic agent allows sensitive and stringent hybridization of short oligonucleotide probes at room temperature [Van Ness and Chen (1991) *Nucl. Acids Res.* 19:5143-5151]. Suitable chaotropic agents include guanidinium chloride, guanidinium thiocyanate, sodium thiocyanate, lithium tetrachloroacetate, sodium perchlorate, rubidium tetrachloroacetate, potassium iodide, and cesium trifluoroacetate, among others. Typically, the chaotropic agent will be present at a final concentration of about 3 M. If desired, one can add formamide to the hybridization mixture, typically 30-50% (v/v).

Various hybridization solutions can be employed. Typically, these comprise from about 20 to 60% volume, preferably 30%, of a polar organic solvent. A common hybridization solution employs about 30-50% v/v formamide, about 0.15 to 1 M sodium chloride, about 0.05 to 0.1 M buffers, such as sodium citrate, Tris-HCl, PIPES or HEPES (pH range about 6-9), about 0.05 to 0.2% detergent, such as sodium dodecylsulfate, or between 0.5-20 mM EDTA, FICOLL (Pharmacia Inc.) (about 300-500 kilodaltons), polyvinylpyrrolidone (about 250-500 kdal), and serum albumin. Also included in the typical hybridization solution will be unlabeled carrier nucleic acids from about 0.1 to 5 mg/mL, fragmented nucleic DNA, e.g., calf thymus or salmon sperm DNA, or yeast RNA, and optionally from about 0.5 to 2% wt./vol. glycine. Other additives may also be included, such as volume exclusion agents which include a variety of polar water-soluble or swellable agents, such as polyethylene glycol, anionic polymers such as polyacrylate or polymethylacrylate, and anionic saccharidic polymers, such as dextran sulfate. Methods of optimizing hybridization conditions are well known to those of skill

in the art (see, e.g., Laboratory Techniques in Biochemistry and Molecular Biology, Vol. 24: Hybridization With Nucleic Acid Probes, P. Tijssen, ed. Elsevier, N.Y., (1993)) and Maniatis, *supra*.

5 The basis of gene expression profiling via micro-array technology relies on comparing an organism under a variety of conditions that result in alteration of the genes expressed. Within the context of the present invention a single population of cells was exposed to a variety of stresses that resulted in the alteration of gene expression. Alternatively, the cellular environment may be kept constant and the genotype may be altered. Typical stresses that result in an alteration in gene
10 expression profile will include, but is not limited to conditions altering the growth of a cell or strain, exposure to mutagens, antibiotics, UV light, gamma-rays, x-rays, phage, macrophages, organic chemicals, inorganic chemicals, environmental pollutants, heavy metals, changes in temperature, changes in pH, conditions producing oxidative damage, DNA damage, anaerobiosis, depletion or
15 addition of nutrients, addition of a growth inhibitor, and desiccation. Non-stressed cells are used for generation of "control" arrays and stressed cells are used to generate an "experimental", "stressed" or "induced" arrays.

In an alternate embodiment the present invention provides a method for quantitating the amount of each protein specifying RNA contained within an
20 organism. This is often necessary in gene expression profile analysis because the quantity of transcript produced as well as its fold elevation is needed for quantitative analysis of the cell's physiological state. The method is applicable to both prokaryotic and eukaryotic organisms including for example, cyanobacteria (such as *Rhodobacter* and *Synechocystis*) yeasts (such as *Saccharomyces*,
25 *Zygosaccharomyces*, *Kluyveromyces*, *Candida*, *Hansenula*, *Debaryomyces*, *Mucor*, *Pichia* and *Torulopsis*), filamentous fungi (such as *Aspergillus* and *Arthrobotrys*), plant cells and animal cells. The method proceeds by generating a comprehensive micro-array as described above, from either total or poly-adenylated RNA, depending on the whether the organism is prokaryotic or
30 eukaryotic. Following the generation of the array, a set of labeled DNA and a set of labeled cDNA are synthesized having complementarity to the ORF's of the array. The signals generated from the independent hybridization of either the labeled DNA or cDNA are used to quantitate the amount of protein specifying RNA contained within a genome.

35 In another embodiment the invention provides a method for gene expression profiling with a reduced signal to noise ratio. This is accomplished using a dual "label swapping" method and is again applicable to both prokaryotic and eukaryotic genomes. "Label swapping" refers to a system where a set of

probes or cDNA generated from control or experimental conditions are labeled with two different labels and mixed prior hybridization with the array. Two sets of control and experimental probes or cDNA's are generated. One of the control sets is labeled with a first label (i.e. cy3) and the other is labeled with a different second label (i.e. cy5). The two differently labeled sets are mixed and then hybridized with the array. The same process is repeated for the experimental conditions and the resulting control and experimental fluorescent signals are compared. This combination of signals provides (a) additional measure of each transcript level and (b) allows for the canceling of any bias associated with differential incorporation of fluorescently labeled nucleotide into cDNA or the hybridization of that cDNA.

The preferred embodiments of the invention are discussed below.

Bulk *E. coli* RNA was reverse transcribed to prepare hybridization probes. Despite the large amount of stable RNA (ribosomal and transfer RNAs) in the template, hybridization to protein-encoding genes was readily detected.

As shown in Figure 1 with IPTG induction, conditions have been optimized to yield highly reliable data. In Figure 1, basal expression levels were plotted on the ordinate, induced levels on the abscissa. Panel A illustrates the results obtained when two Cy3-labeled probes were hybridized to duplicate whole genome array sets. Panel B represents an experiment in which the Cy5-labeled cDNA copy of control RNA and the Cy3-labeled copy of induced RNA were co-annealed to a single slide set. The RNAs used to generate the results in Panel B were each labeled with the other dye to allow a "reciprocal" hybridization. In Panel C, the resulting data were averaged with the data presented in Panel B to yield the scatter plot depicted in Panel C. A second independent set of RNA samples were isolated, their cDNAs labeled with both dyes and products hybridized in both possible combinations to generate the results depicted in Panel D. Panel E displays the averaged results of the two independent experiments depicted in Panels C and D.

Reciprocal Labeling. When the results of a single hybridization experiment using different slide sets as capture reagents for Cy3-labeled cDNA derived from treated and control cells were plotted in log-log form, *lacZYA* induction above the background was detected (Figure 1A); variation of other genes was also significant as indicated by the width of the points falling along the diagonal of this scatter plot. Improvements were observed by labeling the control sample with Cy5 and the induced sample with Cy3 before hybridizing to a single set of 3 slides (Figure 1B). However, there was a skewing of the data away from the abscissa and towards the ordinate (y-axis; Cy5-labeled probe). Averaging of these results

with others obtained using reciprocal copying of the same RNA samples (induced RNA reverse transcribed with Cy5 and control RNA with Cy3) resulted in a decreased variation between the treated and control samples (Figure 1C). Such “label swapping” lessened the skewing and decreased the scatter. The experiment, depicted in Figure 1C, was replicated; fresh cultures were induced and nucleic acids processed to yield the data depicted in Figure 1D. The experiments shown in Figures 1C and 1D each represent four measurements of individual transcript abundance; this repetition and averaging yielded the tight constellation shown in Figure 1E which combined the data of Figure 1C and 1D. Nonetheless, the scatter plot resulting from an experiment using the optimized protocol (Figure 1E) illustrated that measurements of gene expression were still subject to considerable variation when the signal was in the lowest part of the detectable range.

The effect of 1 mM IPTG upon expression of the arrayed genes was investigated. Duplicate RNA preparations of the control and IPTG treated cells were each labeled with Cy3 and Cy5 by first strand cDNA synthesis. Averaging of measurements gave an optimal reliability of the data (Figure 1). Examination of the extent of hybridization to any individual gene revealed a wide dynamic range with more than a thousand fold variation in signal intensity between genes (see Figure 1). The expression of only 8 genes increased by a factor of more than 2 after exposure to 1 mM IPTG for 15 min (Figure 1E). These induced genes are listed in Table 1. Two-fold or greater repression was not observed after this treatment. The most highly induced RNAs corresponded to the *lac* operon structural genes. Examples of the induced genes are *b0956*, *mela*, *uxaA* and *b1783*.

Signal Quantitation. The present invention was applied to monitor the effects of growth stage and medium on gene expression. For these embodiments, signal quantitation was important. The percentage of RNA that programs protein synthesis has been determined under a wide variety of growth regimes (Bremer and Dennis, *Escherichia coli* and *Salmonella: Cellular and Molecular Biology* ASM Press: 922-937 (1996)). The fraction of those protein-specifying transcripts devoted to each arrayed gene was estimated. Hybridization signals arising from annealing of RNA-derived Cy3-labeled cDNA populations were quantitated by dividing by the signal generated using Cy3 fluorescent DNA arising from copying of sheared *E. coli* genomic DNA as a probe. The probe synthesized by copying genomic DNA was used to approximate equimolar transcription of the entire genome. This quantitation allowed calculation of mRNA inventories. Three RNA samples were measured. The samples were isolated from cells growing exponentially in rich medium, from cells growing exponentially in minimal

medium, and from cells in minimal medium transitioning from exponential to stationary phase. RNAs from certain central metabolic (*gapA*, *ptsH*), defense (*ahpC*, *cspC*), DNA metabolic (*hns*), surface structure (*acpP*, *ompACFT*, *lpp*), translation (*rplBCKLMPWX*, *rpmBCI*, *rpsACDHJNS*, *trmD*, *fusA*, *infC*, *tufAB*),
 5 transcription (*rpoAB*), and unassigned (*b4243*) genes (Riley and Labedan, *Escherichia coli and Salmonella: Cellular and Molecular Biology*, ASM press: 2118-2202 (1996)) were abundant (>0.1%, among the top 100 transcripts) in all three samples.

The most highly transcribed genes in actively growing broth-cultured cells
 10 often encoded proteins involved in translation. In contrast, cultures at a similar growth stage in glucose minimal medium, expressed to a very high level several small molecule biosynthetic genes and the means to utilize glucose. Thus, an agreement between these molecular analyses and the accumulated understanding of *E. coli* physiology was observed (*Escherichia coli and Salmonella: Cellular*
 15 *and Molecular Biology*, ASM press). This agreement was underscored in the analysis of cells transitioning from the exponential growth phase; the elevated expression of several *rpoS*-controlled genes corresponded to expectations (*Escherichia coli and Salmonella: Cellular and Molecular Biology*, ASM press).

The genes, each representing between 0.0007% and 1% of the hybridizing
 20 signal, were expressed in LB grown cells. The distribution of genes as a function of expression level is plotted in Figure 2. Figure 3 depicts fractional expression as a function of summed genes with genes ranked by expression level. In Figure 2, the histogram plots the number of genes as a function of expression range. Diagonally striped, solid, and horizontally striped bars reflect distributions
 25 observed in RNAs derived from cells growing exponentially in minimal medium, cells transitioning to stationary phase in minimal medium, and cells growing exponentially in rich medium, respectively. In Figure 3, the fraction (summed open reading frame transcripts/total open reading frame transcripts) was plotted as a function of genes summed. The order in which genes were summed was based
 30 upon expression level with the most highly expressed gene summed first.

Fewer genes were expressed in LB than in minimal medium (Figure 2); the fraction of rare transcripts appeared under-represented in LB medium (Figure 3). The fifty most highly expressed genes in broth-grown cells are listed in left-most columns of Table 2; twenty-six of these intensely transcribed genes encode
 35 proteins involved in translation while three encode chaperones.

The broad distribution analyses (Figures 2 and 3) readily revealed the significant differences observed in expression of *E. coli* when grown in defined and rich media. In minimal media many more genes were transcribed over a

somewhat broader range. The 50 genes most highly expressed in exponentially growing cells cultured in minimal medium with glucose as a carbon/energy source are listed in the middle columns of Table 2. Eight biosynthetic genes were highly expressed (Table 2). Notable among them were *metE*, encoding the aerobic methionine synthase, and *ilvC*, an isoleucine-valine biosynthetic gene subject to feed-forward transcriptional activation (Umbarger, H.E, *Escherichia coli and Salmonella: Cellular and Molecular Biology*, ASM Press (1996)) by its substrates. Both the *ilvC*-encoded enzyme (Petersen et al., *Nucleic Acids Res.* 14:9631-9651 (1986)) and *metE*-encoded enzyme (Green, R. C., *Escherichia coli and Salmonella: Cellular and Molecular Biology*, ASM Press (1996)) are sluggish catalysts. The *metE* product accounts for about 5% of *E. coli* protein when cells are cultured in minimal medium with glucose as a carbon/energy source (VanBogelen et al., *Escherichia coli and Salmonella: Cellular and Molecular Biology*, ASM Press (1996)). Other highly expressed biosynthetic genes included *folE* and *cysK*; the *folE* product, GTP cyclohydrolase I catalyzes both cleavage of the 5-membered ring of guanine and the rearrangement of the ribose moiety of the substrate, GTP (Green et al., *Escherichia coli and Salmonella: Cellular and Molecular Biology*, ASM Press (1996)). *cysK*, encoding *o*-acetylserine(thiol)-lyase isozyme A, is responsible for more than 90% of sulfur fixation under aerobic conditions (Kredich, N. M., *Molecular Biology*, ASM press (1996)). Transcripts of the *pyrBI* operon encoding aspartate transcarbamylase also were highly expressed during exponential growth in minimal medium relative to a broth-grown culture. This expression level is a characteristic signature of strain MG1655 whose aspartate transcarbamylase content is elevated more than 100 fold when grown in the absence of uracil due to an *rph* mutation that is polar on *pyrE* (Jensen, K. F., *J. Bacteriol.* 181:3525-3535 (1993)). The other highly expressed transcripts, *thrL* and *pheF*, encoded, respectively, the threonine leader polypeptide (Landick et al., *Escherichia coli and Salmonella: Cellular and Molecular Biology*, ASM Press (1996)) and the phenylalanine-inhibited first enzyme of the common aromatic pathway. The *pheF* product, one of three isozymes, is estimated to account for more than 80% of the activity catalyzing the first common step of aromatic amino acid synthesis (Pittard, A. J., *Escherichia coli and Salmonella: Cellular and Molecular Biology*, ASM Press (1996)).

In this embodiment, expression of several genes catalyzing fueling reactions was also elevated. Unexpectedly, *aceAB*, encoding the glyoxylate shunt enzymes malate synthase and isocitrate lyase (Cronan and Laporte, *Escherichia coli and Salmonella: Cellular and Molecular Biology*, ASM Press (1996)), was highly expressed. Perhaps the TCA cycle functions in its branched state during

this phase of growth requiring the glyoxylate shunt for anapleurotic replenishment (Neidhardt et al., *Physiology of the Bacterial Cell: A Molecular Approach*, Sinauer Associates, Inc. (1990)). As expected, *ptsHI* transcripts encoding phosphotransferase sugar transport common components (Postma et al.,
 5 *Escherichia coli and Salmonella: Cellular and Molecular Biology*, ASM Press (1996)) also accumulated to a very high titer in glucose-minimal medium.

The present invention was applied to monitor the transcripts of cells transitioning from exponential to stationary phase in defined, minimal medium. During this transition, significant changes in gene expression were expected and
 10 observed. Expressed gene levels were from 0.0023 to 1.6%. A total of 1030 genes, of which 110 have a defined role, did not appear to be expressed. In this embodiment, the 50 most highly expressed genes during this transition are listed in the rightmost columns of Table 2. Significantly, several *rpoS*-regulated genes (Hengge-Aronis, *Escherichia coli and Salmonella: Cellular and Molecular*
 15 *Biology*, ASM press, 1497-1512) including *hdeA* (11 fold), *hdeB* (8.9 fold), *dps* (4.4 fold), *gadA* (8.2 fold) and *gadB* (12 fold) (Castanie-Cornet et al., *J. Bacteriol.* 181:3525-3535 (1999)) as well as *rpoS* (2.6 fold) itself became quite highly expressed. Despite this remodeling of transcription, the overall patterns of gene
 20 number as a function of expression level (Figure 2) and fractional expression as a function of ranked gene (Figure 3) were not as distinct as might have been expected in comparison to the patterns observed for RNA extracted from exponentially growing cells.

The observed expression patterns are summarized in Table 3 where gene products were grouped by metabolic function using an established classification
 25 scheme (Riley and Labedan, *Escherichia coli and Salmonella: Cellular and Molecular Biology*, ASM Press (1996)). Exponential growth in minimal medium elevated the amount of pyrimidine and amino acid biosynthetic transcripts. In contrast cofactor and purine transcripts did not appear to accumulate relative to growth in broth. Expression of glyoxylate shunt and miscellaneous glucose
 30 transcripts was also elevated in minimal medium; the seven-fold elevation of glyoxylate shunt transcripts exceeded the average of that observed for amino acid biosynthetic mRNAs. Expression of genes involved in sulfur fixation was also elevated during growth in minimal medium.

The rapid growth observed in LB was reflected in the gene expression
 35 profile, as was the difference in carbon energy/source between glucose and amino acids. LB-grown cultures displayed elevated expression of genes specifying glucogenic enzymes and of genes whose products degrade small molecules. Expression of the ATP and proton motive force generating machinery, elevated by

a factor of about 2, paralleled increased ribosomal protein, aminoacyl-tRNA synthetase and foldase/usher expression.

Changes observed upon entering the transitional period between exponential and stationary phase growth were less dramatic. Nonetheless, elevation of mRNAs specifying gluconogenic, glycolytic, and TCA cycle enzymes was observed as was an increase in transcripts-encoding enzymes responsible for metabolic pool interconversions and for the non-oxidative branch of the hexose monophosphate shunt. The cells also displayed an increased titer of foldase/usher-specifying and global regulatory function transcripts while transitioning between growth phases.

The present invention was used to monitor the change in gene expression when cells overexpressed *sdiA* gene. The *sdiA* is a positive activator of an operon that includes *ftsQAZ*, genes essential for septation.

RNA isolated from broth grown, exponential phase cultures harboring either a single copy (pUC19/RFM443) or many copies (pDEW140/RFM443) of *sdiA* were compared after conversion into fluorescently labeled cDNA by hybridization to individual genes arrayed on glass slides.

Expression of about 9% of the *E. coli* genes was elevated in the strain containing the multicopy *sdiA* plasmid (Table 4). Transcripts of seven genes involved in cell division were raised 2.1 to 11 fold by amplification of *sdiA* as were a large number (about 20) of genes involved in DNA replication, repair, and degradation. Transcript levels of eight genes whose products alter the susceptibility of *E. coli* to drugs were more highly expressed in the strain containing the gene amplification. This genetic configuration also resulted in elevated expression of several lipopolysaccharide biosynthetic genes (*rfa*) as well as open reading frames encoding membrane structural elements.

Expression of several genes of unknown function was also elevated in response to the presence of multiple copies of *sdiA*. The genes whose transcripts were highly (>6 fold) elevated in response to the multicopy *sdiA* plasmid included: *b0135* (6.4 fold, annotated as putative fimbrial-like protein gene), *b0225* (6.4 fold, a gene apparently co-transcribed with *dinJ* since between them there is only a 3 base pair intergenic region), *b0157* (11 fold, encoding a putative malate dehydrogenase), *b0530* (also known as *sfmA* and predicted to specify a fimbrial like protein was elevated 6.5 fold), *b0712* (encoding a putative carboxylase had a 6.4 fold increase in transcript content) and *b1438* (11 fold elevation in expression).

Around 3% of the *E. coli* genes were repressed in a strain harboring the *sdiA* plasmid relative to the control strain containing the vector (Table 5). The

genes involved in chemotaxis, mobility, and flagella biosynthesis were repressed dramatically. Genes for transport of certain carbohydrate substrates and cations (Fe^{++} and K^+), degradation of corresponding carbon compounds, as well as acetate fermentation were repressed. The presence of pDEW140, a pUC19 derivative harboring *sdiA*, resulted in a 30-fold elevation in detectable *sdiA* transcript. Expression of *sdiA* was very low (0.0015 %, the 4212th most abundant transcript) in LB grown *E. coli* MG1655. The increased expression in the plasmid containing strain raised the transcript rank to about 300.

Genes *ddl*, *ftsQ*, *ftsA*, *ftsZ* and *lpxC* are organized in the order mentioned above in the complex *ftsZ* containing operon, and the above genes are transcribed in the same direction starting with *ddl*. Since the *sdiA*-encoded positive activator drives transcription of a mRNA including *ddl*, *ftsQ*, *ftsA*, *ftsZ*, and *lpxC*, increased quantities of RNA hybridizing to these genes were expected. Amplification of *sdiA* due to its presence on a multicopy plasmid elevated expression of *ddl*, *ftsQ*, *ftsA*, *ftsZ* and *lpxC* 4.6, 8.8, 10, 11 and 3.5 fold, respectively, relative to the strain that harbored pUC19 (Table 4).

In the immediate down stream of *sdiA*, there are *yecF*, followed by *uvrY* and *uvrC* gene, respectively. *uvrY* and *uvrC* genes are transcribed in the same direction as *sdiA* and the *yecF* is transcribed in the opposite direction. Unexpectedly, amplification of *sdiA* elevated expression of two genes downstream of *sdiA* was observed. *uvrY* expression was elevated 12 fold while *uvrC* transcription was increased by a factor of 9 (Table 4). These two genes were transcribed in the same direction as *sdiA*. The expression of *yecF* decreased only slightly.

Amplification of *sdiA* caused the expression of 101 genes to fall by a factor of 2 or more. Among them, 44 were involved in motility and chemotaxis. Thirty four genes were down regulated more than five-fold by *sdiA* amplification. Of these, thirty were involved in chemotaxis or motility (*cheW*; *flgB, C, D, E, F, G, H, I, J, K, L, M, N*; *fliA, C, E, F, G, H, J, L, M, N, P, S, T, Z*; *tar* and *tsr*). The master regulator genes *flhC* and *D* controlling flagella operon expression were lowered by only 30-38 %.

The swarming of strains having single or multiple copies of *sdiA* was examined by spotting four single colony isolates of each strain on semi-solid medium. Since almost all the genes involved in flagella biosynthesis, chemotaxis and motility were dramatically repressed in the *sdiA* overexpression strain, loss of mobility of the *sdiA* overexpression strain was predicted. Experiments were carried out to compare the mobility of the two strains. After 8 hr. at 37°C, the strain containing pUC19 had swarmed (diameter = 32 ± 2.5 mm) while that

containing pDEW140 (*sdiA*⁺) had not (diameter = 3.2±0.4 mm). After 23 h the pUC19 containing strain had filled the petri plate while the strain carrying the *sdiA* amplification had significantly swarmed covering about one half of each plate. This partial phenotype could be explained by either (a) plasmid loss
 5 allowing swarming of a revertant (*sdiA*⁺ haploid) population as ampicillin was exhausted from the medium or (b) *sdiA* amplification only partially compromising motility. To distinguish between these possibilities, the site of inoculation and the edge of the swarm after 23 hr were streaked for single colonies to an ampicillin containing LB agar plate. Massive *sdiA*⁺ plasmid loss from cells at the edge of
 10 the swarm was not observed suggesting that the motility defective phenotype was not an absolute one.

If the role of *sdiA* is to stimulate gene expression required for septation, *sdiA* might coordinate expression of the *ftsZ*-containing operon with action at the origin of replication, *oriC*. The two genes immediately flanking *oriC* are *mioC*
 15 and *gidA*. *mioC* is followed by *asnC* and *asnA*, and *gidA* is followed by *gidB*, *atpI* and *atpB*. All of the genes except *asnA* are transcribed in the same direction. *gidA* and *mioC* were over-transcribed relative to the vector-containing control strain. *mioC* transcript content was elevated 7 fold while those of the *gidA* and *gidB* genes were elevated 4 and 2 fold, respectively. This effect was most
 20 localized; adjoining genes were not over-expressed.

Having found enhanced action around *oriC*, it was reasonable to examine the transcript content of genes surrounding the termini of replication when *sdiA* was amplified. There are multiple termini in *E. coli*. The region surrounding *terB* spans minutes 35.3-37.3 (Berlyn et al., *Escherichia coli* and *Salmonella: Cellular*
 25 *and Molecular Biology* ASM Press: 922-937 (1996)) *sdiA* amplification-elevated expression of 12 of the 88 genes in this region more than 3 fold. Transcripts from another 26 genes in the region were elevated by a factor of 1.5 to 3. Unlike the action observed around the terminus, the stimulation seen in the vicinity of *terB* was diffuse. Interestingly, *tau*, encoding the terminus-utilizing factor, was not
 30 over-expressed. Transcription of *gusR*, located at 36.5 minutes, was elevated 8 fold by *sdiA* amplification (Table 4).

acr genes specify sensitivity to acriflavines, molecules that intercalate into double stranded DNA containing monotonic runs of base pairs. Most *acr* mutants display a defect in acridine efflux; moreover they are often pleiotropic being
 35 hypersensitive to a wide variety of chemicals. Thus hyper-expression of these genes in a strain harboring an *sdiA*-bearing multicopy plasmid could lead to mitomycin C expulsion and the observed resistance to this DNA damaging agent. This expectation of *acr* hyper-expression was confirmed. Evidence for elevated

expression of each *acr* operon was found as indicated by the fold expression reported in Table 4.

Elevated transcription of the *gal* operon genes at minute 17 was observed in the strain bearing the *sdiA* amplification. These genes, moderately expressed when strain MG1655 was grown in LB medium (ranks: *galE* 841, *galT* 1512, *galK* 599; Wei and LaRossa, unpublished), were elevated 3.8, 4.9 and 4.1 fold, respectively. Nearby, at minute 16 is the *ybgIJKL-nei* region. *ybg* genes are organized as *ybgF*, *ybgJ*, *ybgK* and *ybgL*, in that order followed by *nei* gene. These genes, transcribed in the same orientation, could constitute an operon since the open reading frames are densely packed, at times overlapping. *sdiA* amplification elevated expression of these genes 5.2, 4.7, 6.4, 3.8 and 8.6 fold, respectively. *nei* encodes an endonuclease responsible for the excision of oxidized pyrimidines in the double helix.

Two linked genes at minute 44, *b1956* and *b1957* were elevated 6.6 and 14 fold by *sdiA* amplification. Similarly, expression of *b2017* and *b2016*, two genes at minute 45 divergently transcribed from and adjacent to the *his* operon, was elevated 3.8 and 3.5 fold, respectively by the presence of the *sdiA*-containing multicopy plasmid.

Mitomycin C (MMC) is a DNA damaging agent. *E. coli* strain, MG1655, was exposed to MMC, and gene expressions were compared in cells that were harvested at 15 and 40 min post exposure. In the cells that were harvested at 15 min, very little SOS response was detected. At the 40 min, expression of 40 genes was elevated greater than 2 fold relative to the control strain. Among the 40, 13 stress response genes were induced (Table 6) more than 2 fold. The SOS genes that were induced by a 40 min exposure to MMC were *recN*, *dinI*, *sulA*, *lexA*, *recA*, *uvrA*, *dinD*, *priC*, *umuC*, *mioC*, *uvrB*, *ruvA*, and *xseA*.

The SOS responsive genes are *lexA*-dependent. In order to determine the gene expression patterns in the presence and the absence of the SOS response, DM800 and DM803 were exposed to MMC for 40 min and the gene expression profiles were compared. DM800 and DM803 harbor *lexA*⁺ and *lexA*^{ind} alleles, respectively. As expected, when exposed to MMC for 40 min, SOS responsive genes were induced greater than 2 fold in DM800 strain. SOS responsive genes, including *lexA*, were not induced in the DM803 strain (Tables 7 and 8). Many genes that were not induced by MMC in DM800 were induced by the DNA damaging agent in DM803. For examples, the expression of the following genes were induced greater than 2 fold in DM803 but not in DM800 (Tables 7 and 8): among the induced genes are those involved with cell division (i.e., *dicB*, *dicC*,

and *sdiA*); chemotaxis and mobility (i.e., *cheW* and *motA*); and the transport of small molecules (i.e., *cycA*, *fadL*, *chaC*, *codB* and *btuC*).

The present invention is not limited to only highly expressed genes for several reasons. First, reproducible expression measurements were obtained over a wide dynamic range (Figure 1E). Second, the data of Figure 3 and Table 1 illustrate that the *lac* operon expression, although low before IPTG induction, was detected suggesting that most transcripts can be readily measured with the described techniques. Analyses of well-characterized "promoter-down" mutants or spiking experiments may be useful in defining the lower limits of expression that can be observed.

EXAMPLES

The present invention is further defined in the following Examples. It should be understood that these Examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

GENERAL METHODS

Standard recombinant DNA and molecular cloning techniques used in the Examples are well known in the art and are described by Sambrook, J., Fritsch, E. F. and Maniatis, T. *Molecular Cloning: A Laboratory Manual*; Cold Spring Harbor Laboratory Press: Cold Spring Harbor, (1989) (Maniatis) and by T. J. Silhavy, M. L. Bennis, and L. W. Enquist, *Experiments with Gene Fusions*, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1984) and by Ausubel, F. M. et al., *Current Protocols in Molecular Biology*, pub. by Greene Publishing Assoc. and Wiley-Interscience (1987).

The meaning of abbreviations is as follows: "hr" means hour(s), "min" means minute(s), "sec" means second(s), "d" means day(s), "mL" means milliliter(s), "μL" means microliter(s), "nL" means nanoliter(s), "μg" means microgram(s), "ng" means nanogram(s), "mM" means millimole(s), "μM" means micromole(s).

Media and Culture Conditions:

Materials and methods suitable for the maintenance and growth of bacterial cultures were found in Experiments in Molecular Genetics (Jeffrey H. Miller), Cold Spring Harbor Laboratory Press (1972), Manual of Methods for General Bacteriology (Phillip Gerhardt, R.G.E. Murray, Ralph N. Costilow, Eugene W. Nester, Willis A. Wood, Noel R. Krieg and G. Briggs Phillips, eds),

pp. 210-213, American Society for Microbiology, Washington, DC. or Thomas D. Brock in Biotechnology: A Textbook of Industrial Microbiology, Second Edition (1989) Sinauer Associates, Inc., Sunderland MA. All reagents and materials used for the growth and maintenance of bacterial cells were obtained from Aldrich
5 Chemicals (Milwaukee, WI), DIFCO Laboratories (Detroit, MI), Gibco/BRL (Gaithersburg, MD), or Sigma Chemical Company (St. Louis, MO) unless otherwise specified.

LB medium contains following per liter of medium: Bacto-tryptone (10 g), Bacto-yeast extract (5 g), and NaCl (10 g).

- 10 Minimal M9 medium contains following per liter of medium: Na_2HPO_4 (6 g), KH_2PO_4 (3 g), NaCl (0.5 g), and NH_4Cl (1 g).

Above media were autoclaved for sterilization then 10 mL of 0.01 M CaCl_2 and 1 mL of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ plus carbon source and other nutrient were added as mentioned in the examples. All additions were pre-sterilized before they
15 were added to the media.

Molecular Biology Techniques:

Restriction enzyme digestions, ligations, transformations, and methods for agarose gel electrophoresis were performed as described in Sambrook, J., et al., Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor
20 Laboratory Press (1989). Polymerase Chain Reactions (PCR) techniques were found in White, B., PCR Protocols: Current Methods and Applications, Volume 15(1993) Humana Press Inc.

EXAMPLE 1

- Example 1 demonstrates genomic DNA amplification and the preparation
25 of the high density DNA array.
- Amplification of 4290 *E. coli* genes Specific primer pairs (available from Sigma Genosys Biotechnologies, The Woodlands, TX) for each protein-specifying gene of *E. coli* were used in two consecutive PCR amplification reactions. Genomic DNA (30 ng) was used as the template in the first round of PCR amplification, and
30 500-fold diluted PCR products served as templates for PCR re-amplification. Duplicate 50 μL scale reactions were performed. The PCR reactions were catalyzed with ExTaqTM polymerase (Panvera, Madison, WI) with the four dNTPs (Pharmacia), present at 0.25 mM and the primers at 0.5 μM . Twenty-five cycles of denaturation at 95°C for 30 sec, annealing at 64°C for 30 sec and
35 polymerization at 72°C for 2 min were conducted. A 2 μL aliquot of each PCR product was sized by electrophoresis through agarose gels. More than 95% of the second round PCR products displayed visible bands of the correct size. Second round PCR reactions devoid of templates and primers were saved to serve as

negative controls for hybridization capture reagents. One third of each second round PCR reaction was purified using 96-well PCR purification kits (Qiagen, Valencia, CA). The eluted DNAs were dried using a vacuum centrifuge.

- Arraying amplified genes. Twenty microliters of 6M Na₂SCN or 50% DMSO was added to each dried DNA sample (≥ 0.1 ng/nL). A generation II DNA spotter (Molecular Dynamics, Sunnyvale, CA) was used to array the samples onto coated glass slides (Amersham Pharmacia Biotech, Arlington Heights, IL). Aliquots of approximately 1 nL from 1536 resuspended PCR products were arrayed in duplicate on each slide; a set of three slides supported all amplified *E. coli* genes. To serve as controls, 76 specific *E. coli* PCR products, 8 amplified genes of *Klebsiella pneumoniae* and 12 plant cDNA clones were also spotted onto each slide. Spotted glass slides, after baking at 80°C for 2 hr., were stored under vacuum in a desiccator at room temperature.

EXAMPLE 2

- Example 2 demonstrates gene expression analysis. *E. coli* mRNA was isolated, fluorescent labeled cDNA was prepared using mRNA as a template, and the labeled cDNA was hybridized to the high density DNA array. The amount of DNA hybridized to DNA array was quantitated and analyzed.

Microbiological Methods.

- E. coli* MG1655 was cultured with aeration in either the minimal medium, M9 (Miller, J. H., *Experiments in Molecular Genetics*, Cold Spring Harbor (1972)), supplemented with 0.4% glucose or in the rich medium, LB (Miller, J. H., *Experiments in Molecular Genetics*, Cold Spring Harbor (1972)), at 37°C. The overnight culture was diluted 250 fold into fresh medium and aerated by shaking at 37°C. Samples of the minimal medium culture were harvested at A₆₀₀=0.40 (exponential phase) and 1.6 (transition to stationary phase) prior to RNA isolation. An IPTG induction (Miller, J. H., *Experiments in Molecular Genetics*, Cold Spring Harbor (1972)) was performed to examine the specificity with which it effects gene expression. A culture grown overnight in LB at 37°C was diluted 250 fold into fresh LB and aerated at 37°C. When the culture achieved an appropriate density (A₆₀₀=0.40), it was split. To one portion was added IPTG to a final concentration of 1 mM; the untreated sample served as a control. Incubation of both samples was continued with aeration at 37°C for another 15 min (A₆₀₀=0.45 for both cultures) before RNA isolation was initiated.
- RNA Isolation. An equivalent volume of shaved ice was added to 50 mL samples which were pelleted immediately in a refrigerated centrifuge by spinning at 10,410 x g for 2 min. Each resultant pellet was resuspended in a mixture containing 100 μ L of Tris HCl (10 mM, pH 8.0) and 350 μ L of β -mercaptoethanol

supplemented RLT buffer [Qiagen RNeasy Mini Kit, Valencia, CA]. The cell suspension was added to a chilled 2 mL screwed-capped microfuge tube containing 100 μ L of 0.1 mm zirconia/silica beads (Biospec Product Inc., Bartlesville, OK). The cells were broken by agitation at room temperature for 25 sec with a Mini-Beadbeater (TM) (Biospec Products Inc., Bartlesville, OK). Debris was pelleted by centrifugation for 3 min at 16,000 x g and 4°C; the resultant supernatant was mixed with 250 μ L of ethanol. This mixture was loaded onto a column from the Qiagen RNeasy Mini Kit. RNA isolation was completed using the protocol supplied with this kit. Incubation for 1 hr. at 37°C in 40 mM Tris pH 8.0, 10 mM NaCl, 6 mM MgCl₂ with RNase free RQ1 DNase (1 unit/ μ L, Promega, Madison, WI) digested any genomic DNA contaminating the RNA preparation. The digestion products were purified by a second passage through the RNeasy protocol (Qiagen, Valencia, CA). The product was eluted from the column in 50 μ L RNase-free water prior to determining sample concentration by an A₂₆₀ reading. RNA preparations were stored frozen at either -20 or -80°C.

Synthesis of fluorescent cDNA from total RNA. Six microgram of RNA template and 12 μ g of random hexamer primers (Operon Technologies, Inc., Alameda, CA) were diluted with double distilled (dd) water to a volume of 22 μ L. Annealing was accomplished by incubation at 70°C for 10 min followed by 10 min at room temperature. In order were added: 8 μ L of 5x SuperScript II reaction buffer (Life Technologies, Inc., Gaithersburg, MD), 4 μ L of 0.1M DTT, 2 μ L of the dNTP mix (2 mM dATP, 2 mM dGTP, 2 mM TTP, 1 mM dCTP), 2 μ L of 0.5 mM Cy3- or Cy5-dCTP (Amersham Pharmacia Biotech, Arlington Heights, IL), and 2 μ L of SuperScript II reverse transcriptase (200 units/mL, Life Technologies Inc., Gaithersburg, MD). DNA synthesis proceeded at 42°C for 2.5 hr. before the reaction was terminated by heating at 94°C for 5 min. Alkaline hydrolysis of the RNA templates was achieved by adding 2 μ L of 5M NaOH followed by incubation at 37°C for 10 min. Hydrolysis was terminated by the sequential addition of 3 μ L of 5M HCl and 5 μ L of 1M Tris-HCl, pH 6.8. The labeled cDNA was purified with a PCR purification kit (Qiagen, Valencia, CA), dried in a speed vacuum and stored at -20°C. Labeling efficiency was monitored using either A₅₅₀, for Cy3 incorporation, or A₆₅₀, for Cy5 labeling, to A₂₆₀ ratios.

Fluorescent labeling of genomic DNA. Genomic DNA, isolated from strain MG1655 (Bachmann, B., *Escherichia coli* and *Salmonella: Cellular and Molecular Biology*, ASM Press (1996)) by standard procedures (Van Dyk and Rosson, *Methods in Molecular Biology: Bioluminescence Methods and Protocols*, Humana Press Inc. (1998)), was nebulized to approximately 2 kb pair fragments. Three microgram of this DNA was mixed with 6 μ g of random hexamers primers

(Operon Technologies, Inc., Alameda, CA) in 33 μ L of dd water. DNA was denatured by heating at 94°C prior to annealing on ice for 10 min. Fluorescent copying of the genomic DNA was accomplished using the Klenow fragment of DNA polymerase I (5 μ g/ μ L, Promega, Madison, WI). To the DNA mixture was added 6 μ L of 10x Klenow buffer (supplied with the enzyme), 3 μ L of the dNTP mix described above, 12 μ L dd H₂O, 3 μ L of 0.5 mM Cy3-dCTP (Amersham Pharmacia Biotech, Arlington Heights, IL), and 3 μ L of the Klenow fragment of DNA polymerase I. After a static, 2.5 h incubation at room temperature, the labeled DNA probe was purified using a PCR purification kit (Qiagen, Valencia, CA) before drying in a speed vacuum.

Hybridization and washing. Spotted slides were placed in isopropanol for 10 min, boiled in dd H₂O for 5 min and dried by passage of ultra-clean N₂ gas prior to pre-hybridization. The prehybridization solution (PHS) was 3.5xSSC (BRL, Life Technologies Inc., Gaithersburg, MD), 0.2% SDS (BRL, Life Technologies Inc., Gaithersburg, MD), 1% bovine serum albumin (BSA, Fraction V, Sigma, St. Louis, MO). The hybridization solution (HS) contained 4 μ L of dd water, 7.5 μ L of 20xSSC, 2.5 μ L of 1% SDS (BRL, Life Technologies Inc., Gaithersburg, MD), 1 μ L of 10 mg/ml Salmon sperm DNA (Sigma, St. Louis, MO) and 15 μ L of formamide (Sigma, St. Louis, MO). The slides were incubated at 60°C for 20 min in PHS. The slides were next rinsed 5 times in dd water at room temperature and twice in isopropanol before drying by the passage of nitrogen. The dried probe was resuspended in the HS and denatured by heating at 94°C for 5 min.

Thirty microliter of the probe-containing HS was applied to a dried, pre-hybridized slide, covered with a cover slip (Corning, Corning, NY), and put into a sealed hybridization chamber containing a small reservoir of water to maintain moisture. Hybridization occurred for approximately 14 h at 35°C. Cover slips were removed in washing buffer I (WB I = 2xSSC, 0.1% SDS) warmed to 35°C prior to incubation for 5 min. Next, the slides were washed sequentially for 5 min in 1xSSC, 0.1% SDS and 0.1xSSC, 0.1% SDS. Slides were then passed through three baths, each passage lasting 2 min, in 0.1xSSC. The slides were dried with a nitrogen gas flow.

Data Collection and Analysis. Hybridization to each slide was quantified with a confocal laser microscope (Molecular Dynamics, Sunnyvale, CA) whose photomultiplier tube was set to 700 volts and 800 volts for obtaining Cy3 and Cy5 signals respectively. The images were analyzed with ArrayVision 4.0 software (Imaging Research, Inc., Ontario, Canada). The fluorescent intensity associated with each spotted gene was reduced by subtracting the fluorescence of an adjoining, non-spotted region of the slide. These readings were exported to a

spreadsheet for further manipulation. The four “no DNA” spots derived from PCR reactions devoid of template were controls used to determine the noise (background signal) level.

5 The 96 genes present on each slide were used as internal controls to quantify signal intensities yielding equivalent readings among the three slides of a whole genome array set. This corrected for slide-to-slide signal variation.

For the IPTG induction experiment, it was presumed that the overall transcriptional pattern did not change significantly. Thus the summed equivalent reading for the entire genome was quantified; analogous quantitation of the
10 underlying equivalent readings allowed calculation of fold induction of each gene’s expression by comparison of such quantified equivalent readings.
RNA abundance. To convert normalized equivalent readings into measures of transcript abundance, a further correction was needed. That correction required the hybridization signal arising from an equimolar concentration of all transcripts.
15 The surrogate for this transcript pool was the fluorescent copy of genomic DNA. Thus, the fluorescent intensities from hybridization with RNA-derived probes were corrected using fluorescent intensities arising from genomic DNA derived probes. Specifically, the abundance of each gene’s transcription product(s) was determined by dividing the normalized equivalent reading of the genomic DNA
20 derived sample into the normalized equivalent reading from the RNA derived sample. The convention of Riley (Riley and Labedan *Escherichia coli* and *Salmonella: Cellular and Molecular Biology ASM Press*, 1996)) was followed in grouping genes into functional sets.

EXAMPLE 3

25 Example 3 demonstrates gene expression profile changes when cell were exposed to IPTG, or grown in different culture media. The results are illustrated in Tables 1,2 and 3 (Listing of Tables) as described above.

IPTG Induction An *E. coli* strain MG1655 was grown overnight in LB at 37°C. The culture was diluted 250 fold into fresh LB and aerated at 37°C. When the
30 culture achieved an appropriate density ($A_{600}=0.40$), it was split into two portions.

To one portion, IPTG was added to a final concentration of 1 mM. The other portion was untreated and served as a control.

Both samples was incubated with aeration at 37°C for another 15 min
35 ($A_{600}=0.45$ for both cultures) before RNA isolation. Gene expression analysis was performed as described in Examples 1 and 2.

Cells were grown in different culture media *E. coli* MG1655 was cultured with aeration overnight in either the minimal medium, M9, supplemented with

- 0.4% glucose or in the rich medium, LB at 37°C. The overnight culture was diluted 250 fold into fresh medium and aerated by shaking at 37°C. Samples of the minimal medium culture were harvested at $A_{600}=0.40$ (exponential phase) and $A_{600}=1.6$ (transition to stationary phase) prior to RNA isolation. The LB culture was harvested at $A_{600}=0.4$ prior to RNA isolation. Gene expression analysis was performed as described in Examples 1 and 2.

EXAMPLE 4

- Example 4 demonstrates gene expression changes and the effect on mobility when *sdiA* gene was overexpressed in *E. coli*. The results are tabulated in Tables 4 and 5 (Listing of Tables) as described above.

The following plasmids and strains were used in this example.

<u>strain or plasmid</u>	<u>genotype</u>
MG1655	<i>rph-1</i>
RFM443	<i>rpsL galK2 lacΔ74</i>
pUC19	Cloning vector
pDEW140	pUC19 + <i>sdiA</i> (EcoRI)

Strains and growth conditions

- Strains of MG1655 (Bachmann, B., *Escherichia coli* and *Salmonella: Cellular and Molecular Biology*, ASM Press (1996)) and RFM443 (Menzel R., *Anal. Biochem.*, 181:40-50 (1989)) have been described.

- pDEW140 was constructed as following: Chromosomal DNA isolated from *E. coli* W3110 was partially digested with restriction enzyme *Sau3A1* and size fractionated on agarose gels. Fractions of two size ranges (average sizes of approximately 2.5 and 4.0 Kbp) were ligated to pBR322 (0.11 pmol) or pUC18 (0.11 pmol) that had previously been digested with restriction enzyme *BamHI* and treated with calf intestinal alkaline phosphatase. The molar ratio of chromosomal DNA to vector in each of the ligation reactions was approximately 0.2:1. The ligation products were used to transform ultracompetent *E. coli* XL2Blue (Stratagene) to ampicillin resistance. Pooled transformants ($>10^5$ for each transformation) were used to isolate plasmid DNA.

- 0.3 ng of the pUC18 library was electro-transformed into RFM443. The MMC resistant clones were selected on LB agar plates supplemented with 100 $\mu\text{g/mL}$ of ampicillin and 6 mg/mL of MMC. Resistant colonies appeared after the incubation at 37°C. The colonies underwent single colony purification on the same medium. Plasmids derived from single colonies were isolated with the Qiagen 96-well turbo plasmid prep kit. These plasmids served as a template

for primer-directed DNA sequencing of the insert ends. One of the plasmids, Plasmid p[3+4/B10], was shown by sequencing to carry the *sdiA* and surrounding genes. From this plasmid *sdiA* was amplified by PCR using the primers:

f primer = TGGCA CGCAG GACAG AA (SEQ ID NO:1)

5 d primer = TAACA AATCA GCATA ACTCA T (SEQ ID NO:2)

The PCR used Ampli-Taq Gold. Conditions were 94°C, 11 min followed by 32 cycles of 94°C for 45 sec, 45°C for 45 sec, 72°C for 90 sec, the 72°C for 7 min.

10 The PCR product was blunt end ligated into *EcoRV* digested pT7Blue-3 (Novagen). A clone having the proper sized fragment was obtained after transformation into DH5-alpha. From colonies, inserts of the proper size were detected by PCR-based analysis. Such colonies served as a source of plasmid DNA from which *sdiA* was liberated by digestion with *EcoRI*. The fragment was sized by electrophoresis through agarose gels and ligated into *EcoRI* digested
15 pUC19. The ligation mixture was used to transform DH5alpha. Plasmid preps of the transformants were sequenced. One such plasmid containing *sdiA* was named pDEW140 and transformed into strain RFM443.

Plasmids pUC19 and pDEW140 were transformed into RFM 443 selecting for ampicillin resistance on solidified LB agar medium.

20 Strains of RFM443 (pUC19) and RFM443 (pDEW140) were grown overnight with aeration in LB with 150 µg/mL ampicillin (LB with amp). The overnight culture was diluted 250 fold into fresh medium (LB with amp) and incubated further at 37°C with shaking. Cells were collected at O.D.600=0.45, and total RNA was purified using Qiagen RNeasy mini.

25 Motility experiment.

A single colony was picked from freshly grown RFM443 (pUC19) or RFM443 (pDEW140) cultured on LB agar (1.2%), and the center of a LB with amp soft agar (0.3%) plate was stabbed. The soft agar plate containing each culture was incubated at 37°C. The diameters of the growth zones of the two
30 strains were measured and compared.

EXAMPLE 5

Example 5 demonstrates the differences in gene expression profile between strains proficient or deficient in their ability to respond to DNA damaging agents. An isogenic pair of strains, differing only in *lexA*, was used to
35 investigate the cell's range of responses to the DNA damaging agent mitomycin C (MMC). The results are tabulated in Tables 6, 7, and 8 (Listing of Tables) as described above.

Strains *E. coli* strain, MG1655, was used to determine the gene expression profile of *E. coli* in response to a MMC challenge. Two isogenic *E. coli* strains (Mount *et al*, *J. Bacteriol.* 112:886-893 (1972)), DM800 (*lexA*⁺), used as control displaying a normal response to DNA damage, and DM803 (*lexA*^{ind}), a strain
 5 unable to mount the predominant "SOS" response to DNA damage, were compared using comprehensive gene expression profiling.

MMC experiment MG1655 cells were grown in LB overnight with aeration. The overnight cultures were diluted 100 fold in LB to final volume of 500 mL and grown at 37°C to exponential phase. 200 mL of culture was treated with MMC to
 10 the final concentration of 250 ng/mL. Another 200 mL of culture were mock treated without MMC for comparison. Cells were harvested at 15 min and 40 min for MG1655 strain. With DM800 and DM803 stains, cells, cultured in an identical manner, were harvested after 40 min exposure. RNA was isolated and gene expression profile was analyzed as shown in Examples 1 and 2. As seen in
 15 Tables 7 and 8, the *lexA* allele has a great influence on the response of cells to MMC. Table 8 shows that the strain deficient in SOS response still response to MMC but in different manner.

EXAMPLE 6

Preparation of a *Synechocystis* sp. PCC6803 cDNA Probes

20 This example describes the construction of *Synechocystis* sp. PCC6803 cDNA probes following growth of the cells in either minimal growth media (control) or minimal media plus UV-B light treatment. The prepared cDNA probes are used to determine gene expression patterns of many genes simultaneously on a *Synechocystis* sp. PCC6803 DNA microarray as described in
 25 Examples 7 and 8 below.

Hybridization of Microarray Slides and Quantitation of Gene Expression

Microarray glass slides were treated with isopropanol for 10 min, boiling double distilled water for 5 min, then treated with blocking buffer (3.5 x SSC, 0.2% SDS, 1% BSA) for 20 min at 60°C, rinsed five times with double distilled
 30 water, then twice with isopropanol, followed by drying under nitrogen. Cy3 labeled cDNA probes prepared from the total RNA of the UV-B treated *Synechocystis* culture, mixed with an equal amount of Cy5 labeled cDNA probes prepared from the total RNA of the untreated *Synechocystis* culture, were applied to the glass slide in a total volume of 30 µL. The hybridization was repeated
 35 using Cy5 labeled cDNA probes prepared from total RNA of UV-B treated *Synechocystis* culture mixed with an equal amount of Cy3 labeled cDNA probes prepared from the total RNA of the untreated culture, and applied to a second glass slide in a total volume of 30 µL. The hybridization reactions on the glass

slides were performed for 16 hr at 42°C, in a humidified chamber. Hybridized slides were washed in 1X SSC (0.15 M NaCl, 0.015 M sodium citrate), 0.1% SDS for 5 min at 42°C; 0.1X SSC, 0.1% SDS for 5 min at 42°C; three washes in 0.1X SSC for 2 min at room temperature; rinsed with double distilled water and isopropanol; and dried under nitrogen. The slides were scanned using a Molecular Dynamics laser scanner for imaging of Cy3 and Cy5 labeled cDNA probes. The images were analyzed using Array Vision Software (Molecular Dynamics, Imaging Research) to obtain fluorescence signal intensities of each spot (each ORF on the array) to quantitate gene expression. The ratio between the signals in the two channels (red:green) is calculated and the relative intensity of Cy5/Cy3 probes for each spot represents the relative abundance of specific mRNAs in each sample.

Synechocystis Strain and Culture Methods

Briefly, *Synechocystis sp.* PCC6803 cells were grown at 30 $\mu\text{ES}^{-1}\text{m}^{-2}$ light intensity in a minimal growth media, BG-11 (Catalog # C-3061, Sigma Chemical Co., St. Louis, MO) at 30°C, with shaking at 100 rpm with 5% CO₂. Fifty milliliters of *Synechocystis* cells grown to mid logarithmic phase (OD_{730nm} = 0.8 to 1.0) were divided into two 25 mL cultures and transferred from the Erlenmeyer growth flask to two 100 mL petri dishes. The petri dishes, with the lids on, were placed on a rotary shaker and shaken at 100 rpm.

Cell Treatments

For the control, the petri dishes comprising the *Synechocystis* cells were placed on a rotary shaker with the lids on, and shaken at 100 rpm. For the UV-B treated group, the petri dishes comprising the *Synechocystis* cells were placed on a rotary shaker with the lids on, and shaken at 100 rpm. A UV-B lamp (302 nm,) was positioned above the petri dishes and the distance between the UV-B light source and the petri dishes was adjusted to give the desired level of UV-B light intensity. The level of UV-B light intensity was measured at the surface of the cell culture using a UV light meter, following the manufacturer's instructions. UV-B treatment was performed for either 20 min or 120 min. Following UV-B irradiation, the cells were immediately cooled on ice and their RNA isolated as described below.

Total RNA Isolation and cDNA Probe Synthesis

Control-treated *Synechocystis* cells and UV-B treated *Synechocystis* cells were cooled rapidly on ice and centrifuged at 4000 rpm for 5 min. Total RNA samples were isolated using Qiagen RNeasy Mini Kit (Qiagen), following the manufacturer's protocol. RNase A digestion was performed as described in the

protocol, and a second round purification was performed using the RNeasy Mini Kit. The purified total RNA was analyzed by agarose gel electrophoresis.

From each total RNA preparation, both Cy3 and Cy5 fluorescent dye labeled cDNA probes were prepared. To synthesize the Cy3 or Cy5 labeled cDNA probes, a reverse transcription reaction was performed using 10 µg total RNA, 12 µg random hexamer (Ambion), 50 µM of dATP, dGTP, dTTP, 25 µM of dCTP, and 15 µM Cy3-dCTP or 22 µM Cy5-dCTP (Amersham Pharmacia Biotech), DTT, and AMV reverse transcriptase (Gibco BRL). The reaction was carried out at 42°C for 2.5 hr. After the labeling reaction, RNA templates were degraded by alkaline hydrolysis and the cDNA probes were purified using Qiagen PCR purification kit. The purified probes were quantitated by measuring the absorbance at 260 nm, 550 nm (Cy5 dye incorporation) and 650 nm (Cy3 dye incorporation). Prior to hybridization, 100-200 pmol of the purified Cy3 or Cy5 labeled cDNA probes were dried under vacuum, and re-dissolved in the hybridization buffer (5x SSC, 50% formamide, 0.1% SDS, and 0.03 mg/mL salmon sperm DNA).

EXAMPLE 7

Analysis of *Synechocystis sp.* PCC6803 Gene Expression in Minimal Media

Using a *Synechocystis sp.* PCC6803 DNA microarray prepared according to the methods described above and the cDNA probes prepared as described in Example 6, Applicants have identified herein promoters that can be employed for engineering high levels of gene expression in *Synechocystis sp.* PCC6803, other *Synechocystis* species, *Synechococcus*, and like organisms. This Example describes the identification of the most highly expressed genes and their corresponding strong promoters in *Synechocystis sp.* PCC6803 when grown in BG11 media containing 5 mM glucose as described above.

Specifically, a DNA microarray was prepared according to the methods described above using DNA isolated from *Synechocystis sp.* PCC6803 cells grown in BG11 media containing 5 mM glucose. Minimal media *Synechocystis sp.* PCC6803 gene expression was determined by hybridizing this DNA microarray as described above with fluorescent cDNA probes synthesized from total RNA isolated from *Synechocystis sp.* PCC6803 cells grown in BG11 media containing 5mM glucose as described in Example 6.

Briefly, for each minimal media experiment, two hybridization reactions were performed as described above. Specifically, the first reaction used equal molar (typically 100-200 pmol) of Cy5-labeled cDNA from total RNA of the minimal media treated sample, and Cy3-labeled cDNA probes synthesized from *Synechocystis sp.* PCC6803 genomic DNA; the second reaction used Cy3-labeled

cDNA from total RNA of the minimal media treated sample, and Cy5-labeled cDNA probes synthesized from *Synechocystis sp.* PCC6803 genomic DNA. The signal intensities were quantitated as described above. To calculate the ratio of fold induction (i.e., minimal media/genomic), the minimal media treated sample signal intensities were divided by the signal intensities of the genomic sample. As there were two sets of data from duplicated spotting within each slide, the total number of gene expression measurements for each gene was four. All four induction ratios for each gene were analyzed using an Excel program (Microsoft) to determine the standard deviation; an indicator of the level of confidence for the specific data set for each gene. The ratio of signal intensities represents a relative transcription level of each gene in the same experiment. Herein, Applicants have identified the most highly expressed genes, i.e., those genes that are under the control of the strongest promoters, in *Synechocystis* under this minimal media condition (see Table 9).

EXAMPLE 8

Analysis of *Synechocystis sp.* PCC6803 Gene Expression Following UV-B Exposure

Using a *Synechocystis sp.* PCC6803 DNA microarray prepared according to the methods described above and the probes prepared as described above in Example 6, Applicants have identified herein UV-B inducible promoters that can be employed for engineering high levels of gene expression in *Synechocystis sp.* PCC6803, other *Synechocystis* species, *Synechococcus*, and like organisms. This Example describes the identification of the most highly UV-B responsive genes in *Synechocystis sp.* PCC6803 when grown under minimal media conditions and exposed to 20 minutes of UV-B irradiation at $20 \mu\text{ES}^{-1}\text{m}^{-2}$ intensity. These UV inducible promoters can be used to control expression of certain proteins that may be toxic to *Synechocystis* cells.

Specifically, a DNA microarray was prepared according to the methods described above using DNA isolated from *Synechocystis sp.* PCC6803. For each UV-B treatment experiment, two hybridization reactions were performed as described above. In particular, the first reaction used equal molar (typically 100-200 pmol) of Cy5-labeled cDNA from total RNA of the UV-B treated sample, and Cy3-labeled cDNA from total RNA of the control sample (*Synechocystis sp.* PCC6803 grown in BG11 media containing 5 mM glucose); the second reaction used Cy3-labeled cDNA from total RNA of the UV-B treated sample, and Cy5-labeled cDNA from total RNA of the control sample. The signal intensities were quantitated as described above. To calculate the ratio of fold induction (i.e., UV-B/control), the UV-B treated sample signal intensities were

divided by the signal intensities of the control sample. As there were two sets of data from duplicated spotting within each slide, the total number of gene expression measurements for each gene was four. All four induction ratios for each gene were analyzed using an Excel program (Microsoft) to determine the standard deviation; an indicator of the level of confidence for the specific data set for each gene.

Applicants have identified herein the most highly UV-B induced genes in *Synechocystis* following UV-B treatment (see Table 10). Only genes whose expression was induced more than 4 folds by UV-B light (20 min at $20 \mu\text{ES}^{-1}\text{m}^{-2}$ intensity) as compared to the minimal media control are listed in Table 10. The promoters of these genes can be used to construct UV inducible expression vectors in *Synechocystis*.

Some of the gene families induced by UV-B light include D1 protein (psbA), phycobilisome degradation proteins (nblA, nblB), carotenoid biosynthesis enzymes (crtD, crtD, crtQ), chaperones (clpB, ctpA, dnaJ, dnaK, htpG, hsp17), RNA polymerase sigma factor (rpoD), superoxide dismutase (sodB), high light inducible protein (hliA), FtsH protease, which is responsible for the degradation of photo-damaged D1 protein (ftsH), and DNA repair enzyme (uvrC). Among the group of UV inducible genes, there are several genes of unknown function: ssr2016, and sl10185. Applicants' discovery has lead to the first level of functional assignment for these genes. The promoters of these genes can be used to construct UV inducible expression vectors in *Synechocystis*.

A subgroup of Applicants' identified UV-B induced genes comprise two *Escherichia coli*-like -35 promoter sequences in the 5' upstream untranslated regions (UTR), including slr1604 (ftsH), slr0228 (ftsH), sl11867 (psbA3), slr1311 (psbA2), ssl0452 (nblA), ssl0453 (nblA), ssl2542 (hliA), ssr2016 (unknown protein with homologues in green algae and plant), and sl10185 (unknown protein). The nucleotide sequence "GTTACA" is present in the 5' untranslated regions of psbA2, psbA3, and ssr2016 nucleic acids. The nucleotide sequence "TTTACA" was also found to be present in the 5' UTR regions of psbA2, psbA3, ssr2016, rpoD, and ndhD2 nucleic acids.

LISTING OF TABLESTable 1

gene	function	<u>LB</u>	<u>MM exp. phase</u>		<u>MM transition phase</u>	
		fold IPTG induction	fn ^a	rank ^b	fn	rank
<i>lacA</i>	thiogalactoside acyltransferase	36	4.00E-05	3747	2.46E-07	4244
<i>lacZ</i>	galactosidase	29	8.88E-05	2420	2.16E-05	3879
<i>lacY</i>	galactoside permease	14	6.07E-05	3125	6.10E-06	4202
<i>b2324</i>	peptidase?	5.3	2.78E-04	621	7.30E-05	2639
<i>uxaA</i>	altronate hydrolase	4.0	2.93E-04	575	7.82E-05	2530
<i>b1783/yeaG</i>		3.6	3.71E-04	401	3.23E-04	576
<i>melA</i>	galactosidase	2.9	4.05E-05	3729	1.36E-05	4050
<i>b0956/ycbG</i>	hydrogenase?	2.5	2.63E-04	678	1.41E-04	1573

- 5 ^a = fraction of particular transcript/summed transcripts hybridizing to all open reading frames on the micro-arrays;
 ^b = genes are ranked in order of expression with 1 being the most highly expressed gene
 MM: Minimal media,
 exp phase: exponential growth phase

10

Table 2. Highly Expressed Genes under Three Different Culture Conditions

<i>name</i>	fraction ^a in LB	<i>name</i>	fraction in minimal (exp. phase)	<i>name</i>	fraction in minimal (transition)
<i>infC</i> ^b	0.0070	<i>cspA</i>	0.0054	<i>hdeA</i>	0.016
<i>rplK</i>	0.0068	<i>metE</i>	0.0050	<i>hdeB</i>	0.0099
<i>rplL</i>	0.0066	<i>tufB</i>	0.0048	<i>rmf</i>	0.0083
<i>rplA</i>	0.0048	<i>ompA</i>	0.0046	<i>dps</i>	0.0065
<i>hemK</i>	0.0047	<i>ilvC</i>	0.0042	<i>lpp</i>	0.0063
<i>rpmI</i>	0.0046	<i>rmf</i>	0.0038	<i>ompC</i>	0.0059
<i>rplW</i>	0.0044	<i>tufA</i>	0.0038	<i>icdA</i>	0.0059
<i>rplJ</i>	0.0043	<i>ompT</i>	0.0037	<i>metE</i>	0.0053
<i>acpP</i>	0.0042	<i>infC</i>	0.0036	<i>gapA</i>	0.0049
<i>lpp</i>	0.0040	<i>ahpC</i>	0.0034	<i>tufA</i>	0.0049
<i>glpQ</i>	0.0039	<i>rplM</i>	0.0031	<i>ompA</i>	0.0044
<i>fusA</i>	0.0039	<i>ptsH</i>	0.0031	<i>infC</i>	0.0044
<i>gatB</i>	0.0038	<i>aceB</i>	0.0030	<i>uspA</i>	0.0040
<i>rpsF</i>	0.0038	<i>lpp</i>	0.0029	<i>tufB</i>	0.0039
<i>tufB</i>	0.0037	<i>rpsJ</i>	0.0028	<i>ilvC</i>	0.0037
<i>ompC</i>	0.0037	<i>cirA</i>	0.0028	<i>rpsN</i>	0.0036
<i>mopB</i>	0.0035	<i>gapA</i>	0.0026	<i>eno</i>	0.0036
<i>atpF</i>	0.0035	<i>rpmI</i>	0.00266	<i>ahpC</i>	0.0035
<i>hns</i>	0.0035	<i>yjiS</i>	0.0026	<i>ompT</i>	0.0033
<i>rpmB</i>	0.0034	<i>rpmC</i>	0.0024	<i>gadA</i>	0.0033

<i>ompA</i>	0.0033	<i>fusA</i>	0.0024	<i>aceB</i>	0.0033
<i>tnaA</i>	0.0033	<i>b2745</i>	0.0024	<i>rplX</i>	0.0032
<i>rpoA</i>	0.0033	<i>ompF</i>	0.0023	<i>fusA</i>	0.0032
<i>trmD</i>	0.0032	<i>cspC</i>	0.0022	<i>ptsH</i>	0.0031
<i>rplI</i>	0.0031	<i>aceA</i>	0.0021	<i>rpsD</i>	0.0029
<i>gapA</i>	0.0030	<i>pyrB</i>	0.0021	<i>b3512</i>	0.0029
<i>rplM</i>	0.0028	<i>rplK</i>	0.0021	<i>gpmA</i>	0.0028
<i>rpmG</i>	0.0028	<i>rpsD</i>	0.0021	<i>metK</i>	0.0028
<i>rpsC</i>	0.0027	<i>cysK</i>	0.0020	<i>rpmC</i>	0.0027
<i>rplT</i>	0.0027	<i>ptsI</i>	0.0020	<i>gadB</i>	0.0027
<i>rplX</i>	0.0027	<i>b1452</i>	0.0019	<i>rpsV</i>	0.0027
<i>priB</i>	0.0026	<i>rpsS</i>	0.0019	<i>cysK</i>	0.0026
<i>ompF</i>	0.0025	<i>fepA</i>	0.0019	<i>rpsJ</i>	0.0025
<i>hupA</i>	0.0025	<i>pyrI</i>	0.0018	<i>rpsH</i>	0.0025
<i>rpsJ</i>	0.0025	<i>aroF</i>	0.0018	<i>rplE</i>	0.0025
<i>rplB</i>	0.0025	<i>rpsH</i>	0.0017	<i>aceA</i>	0.0025
<i>rplU</i>	0.0025	<i>rpsN</i>	0.0017	<i>b2266</i>	0.0023
<i>tig</i>	0.0024	<i>b0805</i>	0.0017	<i>rplM</i>	0.0023
<i>tufA</i>	0.0024	<i>ompC</i>	0.0017	<i>rpsS</i>	0.0023
<i>rplD</i>	0.0024	<i>rpsA</i>	0.0017	<i>nlpD</i>	0.0022
<i>rplC</i>	0.0024	<i>thrL</i>	0.0017	<i>acpP</i>	0.0022
<i>gatA</i>	0.0024	<i>rplX</i>	0.0016	<i>rpmI</i>	0.0022
<i>rpsA</i>	0.0024	<i>rplL</i>	0.0016	<i>rpoS</i>	0.0021
<i>gatY</i>	0.0023	<i>rpsM</i>	0.0016	<i>rpoA</i>	0.0020
<i>rpsS</i>	0.0023	<i>w4148</i>	0.0016	<i>hns</i>	0.0020
<i>ppa</i>	0.0022	<i>rplB</i>	0.0016	<i>b4253</i>	0.0020
<i>gatZ</i>	0.0022	<i>w0793</i>	0.0016	<i>rplB</i>	0.0020
<i>cspE</i>	0.0021	<i>folE</i>	0.0015	<i>b1452</i>	0.0019
<i>cspC</i>	0.0021	<i>lcdA</i>	0.0015	<i>b0817</i>	0.0019
<i>mopA</i>	0.0021	<i>rplW</i>	0.0015	<i>b1003</i>	0.0019

^afraction of transcripts hybridizing to specified gene/summed transcripts hybridizing to all open reading frames on the micro-arrays

bold, double underlined -foldase/usher genes; **bold, underlined** -stress responsive genes; **bold** - central metabolic enzyme-specifying genes; **double underlined** -biosynthetic genes; **dotted underlined** - translation-associated genes; **underlined** -*rpoS* controlled genes

5

Table 3. Summary of three *E. coli* Expression Profiles

	fraction in MM ^a / exp. phase	fraction in MM/ transition phase	fraction in LB/ exp. phase
1. Cell processes			
<u>Cell division-26^c</u>	0.011	0.010	0.010
<u>Chemotaxis, motility</u>			
Chemotaxis and mobility-12	0.0014	0.00068	0.0011
<u>Folding and ushering proteins-7</u>	0.0032	0.0061	0.011
<u>Transport of large molecules</u>			
Protein, peptide secretion-32	0.0082	0.01014	0.010
<u>Transport of small molecules</u>			
Amino acids, amines-49	0.0091	0.0081	0.0068
Anions-20	0.0029	0.0028	0.0023

Carbohydrates, organic acids, alcohols-82	0.020	0.016	0.034
Cations-52	0.012	0.0098	0.0076
Nucleosides, purines, pyrimidines-6	0.0010	0.00090	0.0017
Other-12	0.0021	0.0027	0.0012
2. Elements of external origin:			
<u>Laterally acquired elements</u>	0.024	0.017	0.023
Colicin-related functions-5			
Phage-related functions and prophages-27	0.0055	0.0042	0.0065
Plasmid-related functions-1	0.00017	0.00055	0.00086
Transposon-related functions-34	0.0058	0.0035	0.0038
3. Global functions			
<u>Energy transfer, ATP-proton motive force-9</u>	0.0077	0.0054	0.015
<u>Global regulatory functions-51</u>	0.0176	0.029	0.018
4. Macromolecule metabolism			
<u>Basic proteins</u>			
Basic proteins - synthesis, modification-6	0.0047	0.0048	0.0074
<u>Macromolecule degradation</u>			
Degradation of DNA-23	0.0038	0.0030	0.0031
Degradation of RNA-11	0.0029	0.0015	0.0022
Degradation of polysaccharides-3	0.00056	0.00033	0.00040
Degradation of proteins, peptides, glyco-61	0.00842	0.0093	0.011
<u>Macromolecule synthesis, modification</u>			
DNA - replication, repair, restr./modific'n-89	0.023	0.019	0.031
Lipoprotein-11	0.0041	0.0050	0.0037
Phospholipids-11	0.0020	0.0015	0.0021
polysaccharides - (cytoplasmic)-6	0.0015	0.0016	0.00060
proteins - translation and modification-34	0.029	0.030	0.043
RNA synthesis, modification, DNA transcript'n-27	0.010	0.010	0.015
<u>Macromolecules</u>			
Glycoprotein			
Lipopolysaccharide-13	0.0015	0.0012	0.0018
aa-tRNAs			
Amino acyl tRNA syn; tRNA modific'n-40	0.013	0.013	0.021
5. Metabolism of small molecules			
<u>Amino acid biosynthesis</u>	0.012	0.0093	0.0033
<u>Biosynthesis of cofactors, carriers</u>	0.072	0.069	0.064
<u>Central intermediary metabolism</u>			
2'-Deoxyribonucleotide metabolism-12	0.0034	0.0032	0.0032
Amino sugars-10	0.0012	0.0011	0.0015
Entner-Doudoroff-3	0.00040	0.00034	0.00060
Gluconeogenesis-4	0.00086	0.0012	0.0021
Glyoxylate bypass-5	0.0076	0.0075	0.0012
Misc. glucose metabolism-3	0.00085	0.00050	0.00039
Non-oxidative branch, pentose pwy-8	0.0026	0.0043	0.0043
Nucleotide hydrolysis-2	0.00010	0.00011	0.00027
Nucleotide interconversions-13	0.0041	0.0039	0.002
Phosphorus compounds-17	0.0032	0.0030	0.0022
Polyamine biosynthesis-8	0.0016	0.0013	0.0013
Salvage of nucleosides and nucleotides-18	0.0037	0.0038	0.0054
Sugar-nucleotide biosynthesis, conversions-18	0.0042	0.0034	0.0048

Sulfur metabolism-10	0.0039	0.0025	0.00095
Pool, multipurpose conversions of intermed. Met'-	0.012	0.019	0.021
46			
<u>Degradation of small molecules</u>			
Amines-9	0.0016	0.0010	0.0025
Amino acids-17	0.0022	0.0014	0.0072
Carbon compounds-90	0.014	0.011	0.025
Fatty acids-10	0.0020	0.0017	0.0030
Other-8	0.0015	0.0021	0.00075
<u>Energy metabolism, carbon</u>			
Aerobic respiration-27	0.0077	0.0058	0.012
Anaerobic respiration-80	0.0075	0.0057	0.011
Electron transport-24	0.0032	0.0024	0.0048
Fermentation-21	0.0040	0.0050	0.0044
Glycolysis-18	0.013	0.024	0.015
Oxidative branch, pentose pwy-2			
Pyruvate dehydrogenase-6	0.0046	0.0041	0.0040
TCA cycle-18	0.0089	0.012	0.0093
<u>Fatty acid biosynthesis</u>			
Fatty acid and phosphatidic acid biosynthesis-23	0.0073	0.0094	0.015
<u>Nucleotide synthesis</u>	0.019	0.013	0.010
Purine ribonucleotide	0.011	0.0077	0.0083
biosynthesis-22			
Pyrimidine ribonucleotide	0.0079	0.0049	0.0018
biosynthesis-10			
6. Miscellaneous	0.37	0.37	0.28
<u>Not classified-109</u>	0.022	0.023	0.025
7. Open reading frames			
<u>Unknown proteins-1324</u>	0.35	0.34	0.26
8. Processes			
<u>Adaptation</u>			
Adaptations, atypical conditions-16	0.012	0.0083	0.0074
Osmotic adaptation-14	0.0038	0.0063	0.0026
<u>Protection responses</u>			
Cell killing-3	0.00052	0.00031	0.00031
Detoxification-11	0.0080	0.0083	0.0097
Drug/analog sensitivity-32	0.0042	0.0031	0.0038
9. Structural elements			
<u>Cell envelope</u>			
Inner membrane-4			
Murein sacculus, peptidoglycan-34	0.0095	0.012	0.013
Outer membrane	0.023	0.026	0.020
constituents-17			
Cell exterior constituents-16	0.0037	0.0039	0.0062
Surface polysaccharides & antigens			
Surface structures-57	0.0075	0.0051	0.0052
<u>Ribosome constituents</u>			
Ribosomal and stable RNAs-3			
Ribosomal proteins - synthesis, modificationRiboso-	0.079	0.086	0.15
54			
Ribosomes - maturation and modification-6	0.0056	0.011	0.00066
10. ORFs n t listed-102			

^aMM = Minimal medium, ^bexp. = exponential, ^cthe number following each description is the number of genes summed

Table 4. Gene expression elevated by the presence of a *sdia* multi-copy plasmid.

5

Genes (grouping by function)	Fold induction	Genes (grouping by function)	Fold induction
1. Cell processes		8. Not classified	
Cell division		<i>agal</i>	3.3
<i>ftsA</i>	10.	<i>chpA</i>	3.0
<i>ftsQ</i>	8.8	<i>dinI</i>	2.0
<i>ftsZ</i>	11	<i>dinP</i>	
<i>minC</i>	2.1	<i>envR</i>	2.2
<i>minD</i>	2.7	<i>ppdB</i>	2.9
<i>minE</i>	2.4	<i>sohA</i>	4.9
<i>sdia</i>	30.	<i>sugE</i>	2.2
<i>sulA</i>	2.6	<i>uvrY</i>	11.9
Chemotaxis and motility		9. Open reading frames of unknown functions	
Transport of large molecules		<i>apaG</i>	2.0
Protein, peptide secretion		<i>hdeB</i>	2.4
<i>msyB</i>	2.1	<i>relE</i>	2.1
<i>oppA</i>	2.1	<i>sprT</i>	3.8
<i>sapB</i>	2.2	<i>b0065</i>	3.7
<i>secD</i>	2.5	<i>b0097</i>	2.1
<i>secF</i>	2.4	<i>b0135</i>	6.4
Transport of small molecules		<i>b0137</i>	2.5
Amino acids, amines		<i>b0138</i>	2.1
<i>glnH</i>	4.0	<i>b0141</i>	4.6
<i>glnQ</i>	2.5	<i>b0163</i>	2.8
Carbohydrates, organic acids, alc		<i>b0189</i>	2.3
<i>araE</i>	5.0	<i>b0224</i>	2.1
<i>frvA</i>	3.2	<i>b0225</i>	6.4
<i>frwD</i>	2.1	<i>b0232</i>	3.1
<i>gntU-1</i>	2.0	<i>b0233</i>	2.9
<i>srlB</i>	2.1	<i>b0234</i>	3.1
<i>xylF</i>	3.8	<i>b0245</i>	2.3
Cations		<i>b0269</i>	3.2
<i>bfr</i>	2.0	<i>b0281</i>	2.4
<i>chaA</i>	2.0	<i>b0295</i>	2.8
<i>feoA</i>	5.8	<i>b0300</i>	2.0
<i>sepD</i>	2.1	<i>b0303</i>	4.7
<i>trkG</i>	4.1	<i>b0322</i>	2.5
2. Elements of external origin:		<i>b0404</i>	3.0
Transposon-related functions		<i>b0407</i>	2.9
<i>rhsC</i>	6.3	<i>b0412</i>	2.4

3. Global regulatory functions		<i>b0443</i>	2.2
<i>lon</i>	2.4	<i>b0461</i>	2.1
<i>lrp</i>	2.2	<i>b0498</i>	2.3
<i>lytB?</i>	3.0	<i>b0517</i>	11.2
<i>rpoE</i>	2.4	<i>b0519</i>	2.5
<i>rseA</i>	2.5	<i>b0530</i>	6.5
<i>rseB</i>	2.2	<i>b0534</i>	2.6
4. Macromolecule metabolism		<i>b0567</i>	2.2
Degradation of proteins, peptides		<i>b0625</i>	3.5
<i>htrA</i>	2.3	<i>b0710</i>	5.2
<i>hycI</i>	2.8	<i>b0711</i>	4.7
<i>ptr</i>	2.1	<i>b0712</i>	6.4
Degradation of DNA		<i>b0713</i>	3.8
<i>endA</i>	2.2	<i>b0715</i>	2.2
<i>mcrB</i>	2.1	<i>b0767</i>	3.3
<i>mcrC</i>	3.5	<i>b1023</i>	2.7
<i>recD</i>	2.4	<i>b1024</i>	2.1
<i>uvrC</i>	9.3	<i>b1069</i>	3.8
Macromolecule synthesis, modifi		<i>b1113</i>	3.8
DNA - replication, repair,		<i>b1214</i>	2.3
<i>gidA</i>	4.1	<i>b1321</i>	2.1
<i>gidB</i>	2.3	<i>b1438</i>	11
<i>hupB</i>	4.5	<i>b1451</i>	2.0
<i>mioC</i>	7.0	<i>b1454</i>	2.2
<i>mutH</i>	2.2	<i>b1455</i>	6.4
<i>nei</i>	8.6	<i>b1458</i>	2.3
<i>priC</i>	2.8	<i>b1463</i>	2.1
<i>recN</i>	3.6	<i>b1487</i>	3.0
<i>umuC</i>	2.3	<i>b1491</i>	2.3
<i>uvrA</i>	2.0	<i>b1498</i>	3.7
<i>xerD</i>	2.3	<i>b1499</i>	2.6
Lipoprotein		<i>b1504</i>	2.1
<i>blc</i>	2.8	<i>b1540</i>	2.5
<i>nlpC</i>	3.2	<i>b1541</i>	6.2
<i>vacJ</i>	2.1	<i>b1542</i>	3.2
Phospholipids		<i>b1543</i>	2.4
<i>pgsA</i>	2.4	<i>b1544</i>	3.0
polysaccharides - (cytoplasmic)		<i>b1545</i>	4.4
<i>glgC</i>	2.1	<i>b1547</i>	2.2
<i>glgS</i>	2.1	<i>b1551</i>	2.1
proteins - translation and modifi		<i>b1560</i>	3.8
<i>prfH</i>	2.2	<i>b1565</i>	3.4
Lipopolysaccharide		<i>b1567</i>	3.5
<i>rfaK</i>	2.3	<i>b1568</i>	2.0
<i>rfaL</i>	2.1	<i>b1579</i>	2.6
<i>rfaY</i>	2.1	<i>b1586</i>	2.2

<i>rfaZ</i>	2.5	<i>b1601</i>	4.0
5. Metabolism of small molecule		<i>b1606</i>	8.0
Amino acids		<i>b1607</i>	7.7
<i>argA</i>	3.4	<i>b1624</i>	2.1
<i>aroD</i>	2.6	<i>b1627</i>	2.4
<i>glnA</i>	2.5	<i>b1628</i>	3.0
<i>glnD</i>	2.6	<i>b1632</i>	2.1
<i>lysR</i>	3.6	<i>b1648</i>	2.6
Biosynthesis of cofactors, carrier		<i>b1649</i>	2.7
<i>thiD</i>	2.1	<i>b1657</i>	4.7
<i>thiM</i>	2.2	<i>b1664</i>	2.1
<i>gst</i>	3.1	<i>b1673</i>	2.4
Central intermediary metabolism		<i>b1688</i>	2.6
2'-Deoxyribonucleotide metabolism		<i>b1699</i>	2.0
<i>nrdA</i>	2.1	<i>b1700</i>	3.7
Amino sugars		<i>b1701</i>	2.0
<i>agaD</i>	4.5	<i>b1706</i>	2.4
Gluconeogenesis		<i>b1707</i>	3.0
<i>ppsA</i>	2.4	<i>b1721</i>	6.5
Phosphorus compounds		<i>b1724</i>	2.1
<i>psiF</i>	3.2	<i>b1743</i>	2.2
Polyamine biosynthesis		<i>b1744</i>	2.5
<i>speC</i>	9.5	<i>b1746</i>	2.9
Salvage of nucleosides and nucleotides		<i>b1756</i>	3.2
<i>apt</i>	2.4	<i>b1789</i>	3.6
<i>gsk</i>	2.4	<i>b1847</i>	2.3
Pool, multipurpose conversions of		<i>b1848</i>	2.7
<i>galM</i>	2.3	<i>b1870</i>	2.4
<i>gcvA</i>	4.6	<i>b1871</i>	2.8
<i>glnK</i>	2.1	<i>b1875</i>	3.4
<i>pntA</i>	10.4	<i>b1877</i>	2.4
<i>pntB</i>	8.2	<i>b1935</i>	2.1
Degradation of small molecules		<i>b1953</i>	3.5
Amino acids		<i>b1955</i>	5.1
<i>tdcB</i>	2.0	<i>b1956</i>	1.4
<i>tdcR</i>		<i>b1965</i>	5.2
Carbon compounds		<i>b1967</i>	6.6
<i>fucA</i>	2.8	<i>b1968</i>	2.4
<i>fucU</i>	1.4	<i>b2006</i>	4.4
<i>galE</i>	3.8	<i>b2007</i>	2.1
<i>galK</i>	4.1	<i>b2015</i>	3.6
<i>galT</i>	4.9	<i>b2016</i>	3.5
<i>glcD</i>	2.0	<i>b2017</i>	3.8
<i>gusR (uidR)</i>	8.0	<i>b2061</i>	2.2
<i>lacA</i>	3.7	<i>b2071</i>	3.1
<i>lacI</i>	2.5	<i>b2145</i>	2.7

	<i>uxuC</i>		<i>b2190</i>	2.3
Fatty acids			<i>b2229</i>	2.1
	<i>atoD</i>	3.8	<i>b2247</i>	2.3
Energy metabolism, carbon			<i>b2253</i>	2.3
Aerobic respiration			<i>b2267</i>	2.0
	<i>nuoH</i>	2.1	<i>b2268</i>	2.0
	<i>nuoI</i>	2.1	<i>b2269</i>	2.6
Anaerobic respiration			<i>b2270</i>	2.2
	<i>dniR</i>	5.8	<i>b2301</i>	3.8
	<i>hybD</i>	3.8	<i>b2302</i>	7.5
	<i>hybE</i>	12	<i>b2339</i>	2.0
	<i>hybF</i>	6.6	<i>b2352</i>	2.6
	<i>hycA</i>	3.1	<i>b2356</i>	2.3
	<i>hycG</i>	2.7	<i>b2385</i>	2.1
	<i>hycH</i>	3.8	<i>b2387</i>	3.6
	<i>hydN</i>	13	<i>b2419</i>	3.2
	<i>hypC</i>	2.3	<i>b2420</i>	3.2
	<i>nrfB</i>	2.7	<i>b2439</i>	2.1
	<i>nrfG</i>	2.1	<i>b2443</i>	2.7
Electron transport			<i>b2444</i>	2.1
	<i>appB</i>	2.5	<i>b2445</i>	2.7
	<i>cybC</i>	2.1	<i>b2485</i>	2.2
Pyruvate dehydrogenase			<i>b2505</i>	2.1
	<i>pdhR</i>	4.4	<i>b2597</i>	2.8
TCA cycle			<i>b2628</i>	2.2
	<i>fumC</i>	5.8	<i>b2629</i>	3.4
	<i>sucA</i>	2.1	<i>b2631</i>	2.9
	<i>sucB</i>	2.6	<i>b2632</i>	4.2
	<i>sucC</i>	2.2	<i>b2640</i>	5.0
	<i>sucD</i>	2.7	<i>b2641</i>	2.3
Fatty acid and phosphatidic acid			<i>b2642</i>	15
	<i>aas</i>	2.0	<i>b2643</i>	2.6
	<i>cdh</i>	2.3	<i>b2648</i>	3.7
Purine ribonucleotide biosynthes			<i>b2649</i>	3.0
	<i>purE</i>	2.1	<i>b2756</i>	2.4
	<i>purR</i>	2.3	<i>b2767</i>	2.9
Pyrimidine ribonucleotide			<i>b2833</i>	2.1
	<i>pyrL</i>	2.1	<i>b2845</i>	2.6
6. Processes			<i>b2846</i>	3.0
Detoxification			<i>b2851</i>	2.6
	<i>cutC</i>	2.1	<i>b2862</i>	2.4
Drug/analog sensitivity			<i>b2874</i>	3.3
	<i>acrA</i>	6.8	<i>b2912</i>	2.6
	<i>acrD</i>	3.0	<i>b2931</i>	2.2
	<i>acrE</i>	14	<i>b2984</i>	2.1
	<i>acrF</i>	6.3	<i>b3021</i>	2.4

<i>acrR</i>	4.5	<i>b3022</i>	2.7
<i>ampC</i>	2.6	<i>b3047</i>	2.2
<i>arsC</i>	2.1	<i>b3050</i>	2.2
<i>tolC</i>	2.6	<i>b3130</i>	2.7
7. Structural elements		<i>b3142</i>	--
<u>Cell envelope</u>		<i>b3254</i>	3.8
Inner membrane		<i>b3372</i>	2.2
<i>smpA</i>	2.6	<i>b3379</i>	2.0
Murein sacculus, peptidoglycan		<i>b3395</i>	4.6
<i>ddlB</i>	4.6	<i>b3397</i>	3.5
<i>hipB</i>	2.0	<i>b3398</i>	2.2
<i>mreD</i>	2.0	<i>b3441</i>	4.0
Outer membrane constituents		<i>b3465</i>	2.1
<i>slp</i>	2.5	<i>b3467</i>	2.5
Cell exterior constituents		<i>b3487</i>	2.1
<i>kdsA</i>	2.3	<i>b3494</i>	2.2
<i>lpxC</i>	3.4	<i>b3513</i>	5.3
<i>rfaB?</i>	2.6	<i>b3535</i>	2.2
Ribosome constituents		<i>b3536</i>	2.9
Ribosomal proteins - synthesis,		<i>b3548</i>	2.0
<i>rpsL</i>	3.1	<i>b3615</i>	2.0
		<i>b3697</i>	2.9
		<i>b3711</i>	2.3
		<i>b3712</i>	2.1
		<i>b3713</i>	2.1
		<i>b3714</i>	2.2
		<i>b3719</i>	2.5
		<i>b3720</i>	3.0
		<i>b3776</i>	2.9
		<i>b3820</i>	2.5
		<i>b3888</i>	3.4
		<i>b3937</i>	2.1
		<i>b3944</i>	2.0
		<i>b3964</i>	2.2
		<i>b4038</i>	2.4
		<i>b4068</i>	2.5
		<i>b4141</i>	2.3
		<i>b4156</i>	2.2
		<i>b4191</i>	5.1
		<i>b4221</i>	5.0
		<i>b4222</i>	4.8
		<i>b4234</i>	2.9
		<i>b4248</i>	2.9
		<i>b4282</i>	2.1
		<i>b4298</i>	2.3
		<i>b4300</i>	2.2

<i>b4325</i>	2.1
<i>b2088</i>	9.4
<i>b4404</i>	5.9
<i>b4405</i>	4.2
<i>yieD (b3695)</i>	3.8
<i>b3914</i>	3.5
<i>b3913</i>	3.3
<i>yjiT (b4352)</i>	2.8
<i>yjiU (b4342)</i>	2.7
<i>yjiQ (b4364)</i>	2.4
<i>yhiL (b3486)</i>	2.3
<i>b2848</i>	2.0
<i>b3573</i>	2.0

Table 5. Gene expression reduced by the presence of a *sdiA* multi-copy plasmid.

Genes (grouping by function)	Fold repression	Genes (grouping by function)	Fold repression
1. Cell processes		6. Structural elements	
Cell division		Outer membrane constituents	
<i>ftsX</i>	2.4	<i>flu</i>	7.7
Taxis and mobility		Cell exterior constituents	
<i>air (aer)</i>	4.6	<i>nanA</i>	3.7
<i>cheA</i>	3.7	Surface structures	
<i>cheB</i>	3.3	<i>fimA</i>	2.6
<i>cheR</i>	2.2	<i>flgA</i>	2.3
<i>cheW</i>	5.3	<i>flgB</i>	14
<i>cheY</i>	4.6	<i>flgC</i>	17
<i>cheZ</i>	4.0	<i>flgD</i>	17
<i>motA</i>	2.9	<i>flgE</i>	17
<i>motB</i>	2.9	<i>flgF</i>	7.1
<i>tar</i>	5.3	<i>flgG</i>	13
<i>tsr</i>	5.9	<i>flgH</i>	5.9
Transport		<i>flgI</i>	5.6
Protein, peptide secretion		<i>flgJ</i>	6.3
<i>dppA</i>	2.3	<i>flgK</i>	6.3
Amino acids, amines		<i>flgL</i>	11
<i>sdaC</i>	3.9	<i>flgM</i>	7.1
Carbohydrates, organic acids		<i>flgN</i>	5.6
alcohols		<i>flhA</i>	3.1
<i>fadL</i>	2.5		

	<i>glpF</i>	3.6		<i>fliA</i>	10.
	<i>glpT</i>	2.5		<i>fliC</i>	14.
	<i>lamB</i>	3.0		<i>fliD</i>	4.6
	<i>malE</i>	3.7		<i>fliE</i>	5.9
	<i>mglA</i>	2.7		<i>fliF</i>	11
	<i>rbsA</i>	2.6		<i>fliG</i>	9.1
	<i>treB</i>	5.3		<i>fliH</i>	5.0
Cations				<i>fliI</i>	3.9
	<i>fecA</i>	3.5		<i>fliJ</i>	7.1
	<i>fecB</i>	2.7		<i>fliK</i>	4.8
	<i>fecE</i>	3.2		<i>fliL</i>	7.7
	<i>fin</i>	2.6		<i>fliM</i>	13
	<i>kdpA</i>	3.5		<i>fliN</i>	5.0
2. Elements of external origin:				<i>fliO</i>	4.4
Phage-related functions and prophages				<i>fliP</i>	6.3
	<i>lar</i>	4.0		<i>fliR</i>	3.1
	<i>nmpC</i>	2.1		<i>fliS</i>	5.9
3. Global regulatory functions				<i>fliT</i>	5.6
	<i>cytR</i>	2.2		<i>fliZ</i>	8.3
4. Degradation of macromolec			Ribosomes - maturation and modification		
DNA			<i>gutM</i>	2.1	
	<i>xseB</i>	4.8	7. Not classified		
Proteins, peptides			<i>fsr</i>	2.1	
	<i>pepE</i>	2.6	8. Open reading frames with unknown functions		
5. Metabolism of small molecu			<i>b0105</i>	2.4	
Pool, multipurpose conversions of intermed. Met'			<i>b0235</i>	2.1	
	<i>glpK</i>	3.5	<i>b0290</i>	3.5	
	<i>glpQ</i>	2.3	<i>b0307</i>	2.4	
	<i>gltF</i>	2.6	<i>b0704</i>	2.7	
Degradation of small molecule			<i>b0732</i>	2.0	
Amino acids			<i>b1100</i>	2.5	
	<i>sdaB</i>	3.2	<i>b1194</i>	5.0	
	<i>tnaA</i>	4.0	<i>b1200</i>	2.9	
	<i>tnaL</i>	2.4	<i>b1329</i>	2.2	
Carbon compounds			<i>b1339</i>	2.2	
	<i>fucR</i>	2.8	<i>b1383</i>	2.4	
	<i>lacZ</i>	6.7	<i>b1520</i>	2.7	
	<i>malM</i>	3.0	<i>b1566</i>	4.4	

<i>malT</i>	2.6	<i>b1690</i>	2.1
<i>treC</i>	2.6	<i>b1722</i>	2.2
Energy metabolism, carbon		<i>b1880</i>	3.6
Anaerobic respiration		<i>b1929</i>	2.1
<i>hypD</i>	2.2	<i>b1930</i>	3.1
Fermentation		<i>b2001</i>	2.3
<i>ackA</i>	3.1	<i>b2005</i>	2.0
<i>aldA</i>	3.3	<i>b2014</i>	2.4
<i>pta</i>	2.3	<i>b2537</i>	2.2
Fatty acid and phosphatidic acid biosynth		<i>b2844</i>	2.7
<i>accD</i>	3.0	<i>b3010</i>	2.2
Purine ribonucleotide biosynth		<i>b3111</i>	2.6
<i>ndk</i>	2.2	<i>b3323</i>	2.9
		<i>b3442</i>	2.5
		<i>b3539</i>	2.4
		<i>b3872</i>	2.1
		<i>yjiZ (b4354)</i>	3.3
		<i>yjbP (b3877)</i>	4.0
		<i>yhjH(b3524)</i>	4.8

Table 6. Gene expression profiles of MG1655 strain when exposed to MMC

m15	control	final ratio	Gene name	M40	contr I	total ratio	Gene name
25115.41	4700.39	5.34	recN	98536.33	11895.47	8.28	recN
34003.66	4565.91	7.45	hslS	11287.86	1677.72	6.73	hslS
1146.46	769.58	1.49	dinI	10515.16	1564.72	6.72	dinI
2516.14	1468.51	1.71	sulA	13936.20	2416.81	5.77	sulA
4531.55	2030.86	2.23	w1816	18614.34	3231.00	5.76	w1816
34497.54	4438.78	7.77	hslT	21578.60	5358.30	4.03	hslT
8490.60	4896.22	1.73	lexA	22768.75	6375.09	3.57	lexA
992.44	1322.83	0.75	w0522	8750.63	2628.39	3.33	w0522
3207.65	2219.58	1.45	w1815	16206.30	4874.54	3.32	w1815
1844.00	1864.81	0.99	w2141	2156.14	691.51	3.12	w2141
21169.76	11055.79	1.91	recA	34375.81	11386.61	3.02	recA
756.88	510.08	1.48	smpA	1614.58	558.50	2.89	smpA
17140.20	23337.13	0.73	cspA	12566.23	4347.87	2.89	cspA
165.71	396.95	0.42	ecpD	227.53	80.37	2.83	ecpD
780.07	941.22	0.83	w3019	802.62	288.87	2.78	w3019
821.35	733.39	1.12	w2878	1151.78	439.75	2.62	w2878
301.47	384.64	0.78	entD	491.14	189.39	2.59	entD
385.41	778.62	0.49	fhuC	818.05	321.73	2.54	fhuC
7203.63	6899.86	1.04	w1201	4623.01	1874.23	2.47	w1201
1476.06	1291.17	1.14	w2999	1710.57	709.13	2.41	w2999
269.15	393.30	0.68	caiB	326.51	139.49	2.34	caiB
5327.31	6525.35	0.82	infA	8186.35	3504.89	2.34	infA
9150.34	6624.37	1.38	uvrA	33530.31	14452.89	2.32	uvrA
1657.22	1816.09	0.91	w2879	3946.52	1727.30	2.28	w2879
4322.77	5547.40	0.78	insB_2	6522.25	2894.35	2.25	insB_2
2310.26	1778.19	1.30	dinD	5316.67	2385.63	2.23	dinD
5349.03	4945.79	1.08	secG	6754.21	3076.60	2.20	secG
136.50	367.57	0.37	priC	341.80	156.11	2.19	priC
617.58	603.54	1.02	w0561	11989.75	5479.59	2.19	w0561
2228.66	2966.21	0.75	exbD	7165.39	3289.27	2.18	exbD
1282.58	893.81	1.43	umuC	5697.94	2659.80	2.14	umuC
6703.55	7422.39	0.90	mioC	8113.11	3804.25	2.13	mioC
3289.24	4228.67	0.78	insB_1	6065.02	2854.20	2.12	insB_1
4042.60	3531.25	1.14	trkH	17795.29	8430.83	2.11	trkH
867.30	1494.75	0.58	w1345	1026.33	487.58	2.10	w1345
541.83	848.15	0.64	dniR	1878.65	899.98	2.09	dniR
5469.65	4392.09	1.25	uvrB	14508.65	6960.78	2.08	uvrB
1561.63	2155.11	0.72	insA_4	2298.22	1111.79	2.07	insA_4
5398.87	3786.47	1.43	ruvA	10492.52	5134.38	2.04	ruvA
343.85	654.22	0.53	appY	815.11	400.28	2.04	appY
18257.00	17197.04	1.06	xseA	13206.64	6494.62	2.03	xseA
1863.91	1771.47	1.05	w0224	6686.92	3310.73	2.02	w0224
5595.42	6241.55	0.90	w3139	11174.79	5555.32	2.01	w3139
1656.98	1560.47	1.06	w2512	24511.01	12207.10	2.01	w2512
349.40	648.41	0.54	w3304	428.93	850.84	0.50	w3304
297.42	326.64	0.91	w2228	271.97	539.98	0.50	w2228
226.05	501.68	0.45	chaB	721.77	1433.44	0.50	chaB
678.56	880.61	0.77	cydA	699.94	1392.46	0.50	cydA
1422.74	2311.13	0.62	melR	5081.98	10140.17	0.50	melR
1051.64	763.06	1.38	w1004	1265.66	2528.73	0.50	w1004
386.58	562.93	0.69	hofG	315.25	630.05	0.50	hofG
513.59	611.47	0.84	w1429	360.81	721.16	0.50	w1429
1256.86	1688.60	0.74	w0299	9464.02	18943.08	0.50	w0299

m15	control	final ratio	Gene name	M40	c ntrol	t tal rati	G ne name
695.83	807.90	0.86	rpiR	748.66	1502.59	0.50	rpiR
904.58	1028.97	0.88	celD	2009.30	4041.66	0.50	celD
2139.22	1712.26	1.25	w0801	2801.71	5636.83	0.50	w0801
749.35	812.05	0.92	w0241	712.18	1434.35	0.50	w0241
827.51	522.31	1.58	w0621	1293.49	2605.36	0.50	w0621
3746.49	3442.41	1.09	putP	12834.08	25859.69	0.50	putP
428.60	231.66	1.85	w4099	377.87	764.43	0.49	w4099
197.66	171.90	1.15	prsA	535.40	1091.60	0.49	prsA
158.38	114.75	1.38	hybD	177.80	363.82	0.49	hybD
83.33	438.61	0.19	sapB	501.32	1027.11	0.49	sapB
334.88	643.56	0.52	w3821	412.40	845.78	0.49	w3821
388.49	568.82	0.68	w1459	417.69	857.67	0.49	w1459
275.47	170.99	1.61	agaD	222.68	459.00	0.49	agaD
309.42	402.34	0.77	ccmD	646.05	1331.95	0.49	ccmD
328.75	360.49	0.91	cpsG	1070.07	2212.14	0.48	cpsG
421.02	702.69	0.60	relB	1002.28	2072.50	0.48	relB
1361.48	1629.16	0.84	w2809	3556.91	7365.32	0.48	w2809
924.92	626.14	1.48	w0824	1242.77	2589.98	0.48	w0824
794.28	965.82	0.82	osmE	2901.61	6072.91	0.48	osmE
277.27	66.53	4.17	w0362	1234.58	2592.98	0.48	w0362
612.37	471.85	1.30	w1927	481.28	1014.05	0.47	w1927
726.43	629.35	1.15	w0211	639.90	1353.69	0.47	w0211
779.65	724.87	1.08	w0237	855.85	1815.70	0.47	w0237
853.25	919.94	0.93	w2592	585.96	1247.13	0.47	w2592
829.44	1006.55	0.82	phnH	987.67	2146.99	0.46	phnH
537.66	1161.05	0.46	flgA	577.37	1255.75	0.46	flgA
656.68	723.92	0.91	w2595	377.12	821.43	0.46	w2595
800.14	800.17	1.00	w2600	792.47	1732.89	0.46	w2600
69.56	216.88	0.32	pheL	93.28	205.15	0.45	pheL
892.86	431.26	2.07	w3049	1146.62	2524.70	0.45	w3049
764.84	318.13	2.40	w1031	1045.43	2305.88	0.45	w1031
937.41	1834.97	0.51	w0295	473.49	1044.97	0.45	w0295
486.50	540.31	0.90	marB	872.85	1928.83	0.45	marB
587.05	622.95	0.94	w0665	786.69	1744.63	0.45	w0665
711.12	397.24	1.79	w1016	1399.58	3109.10	0.45	w1016
903.55	1024.80	0.88	w0298	8569.57	19066.70	0.45	w0298
570.42	464.88	1.23	w0812	1381.42	3073.94	0.45	w0812
1112.52	446.12	2.49	w2026	1456.45	3270.04	0.45	w2026
909.59	321.83	2.83	w0715	1050.65	2361.56	0.44	w0715
739.31	960.38	0.77	pyrL	689.56	1552.58	0.44	pyrL
1533.58	1350.22	1.14	menE	1802.56	4058.69	0.44	menE
38.16	216.63	0.18	rnb	703.05	1594.10	0.44	rnb
707.80	1099.36	0.64	fucR	1770.89	4026.45	0.44	fucR
973.74	681.23	1.43	w2818	278.71	634.22	0.44	w2818
603.99	388.89	1.55	acpD	1990.24	4539.42	0.44	acpD
610.87	662.57	0.92	w0489	332.52	758.67	0.44	w0489
144.96	121.61	1.19	ppdA	177.26	405.05	0.44	ppdA
439.04	478.63	0.92	w1966	488.90	1118.27	0.44	w1966
277.30	324.71	0.85	no template	333.36	764.15	0.44	no template
1245.39	1015.95	1.23	w2401	1085.45	2497.91	0.43	w2401
570.79	872.26	0.65	w4094	1505.90	3477.91	0.43	w4094
100.79	255.85	0.39	dicC	543.43	1266.31	0.43	dicC
1035.07	1108.91	0.93	w0286	844.03	1986.63	0.42	w0286
4510.92	4439.04	1.02	selB	341.91	808.87	0.42	selB
1625.34	1568.78	1.04	w2733	3589.01	8496.95	0.42	w2733

187.58	341.03	0.55	no template	301.58	716.01	0.42	no template
1841.41	2180.01	0.84	trpC	10007.85	23940.58	0.42	trpC
m15	control	final ratio	Gene name	M40	control	total ratio	Gene name
278.39	137.91	2.02	relF	449.81	1077.54	0.42	relF
791.31	342.34	2.31	w1318	1106.98	2655.28	0.42	w1318
224.27	312.05	0.72	agaV	202.28	490.11	0.41	agaV
791.41	366.08	2.16	w1002	1315.80	3189.26	0.41	w1002
890.33	701.73	1.27	w0685	1124.73	2728.61	0.41	w0685
622.65	681.13	0.91	potH	500.10	1228.03	0.41	potH
993.12	584.36	1.70	w2399	853.26	2107.06	0.40	w2399
1275.28	924.58	1.38	metA	991.35	2449.36	0.40	metA
146.54	178.28	0.82	lytB	310.99	770.46	0.40	lytB
419.51	730.52	0.57	w2987	638.04	1593.08	0.40	w2987
827.42	695.16	1.19	w0552	1225.71	3084.40	0.40	w0552
568.15	412.40	1.38	w1846	382.41	968.10	0.40	w1846
114.29	119.19	0.96	dicB	533.34	1351.95	0.39	dicB
842.23	490.95	1.72	w1005	970.38	2465.94	0.39	w1005
912.78	720.90	1.27	w2587	509.45	1301.74	0.39	w2587
1229.46	751.44	1.64	w1260	963.45	2507.31	0.38	w1260
1448.48	1055.79	1.37	w3068	12241.51	31901.37	0.38	w3068
1010.15	949.49	1.06	w0551	2143.72	5619.44	0.38	w0551
794.65	573.71	1.39	w2599	616.94	1633.89	0.38	w2599
905.72	963.26	0.94	w0569	912.64	2444.12	0.37	w0569
1708.81	2679.92	0.64	fruR	7146.34	19274.42	0.37	fruR
1170.58	1168.59	1.00	w3927	1323.78	3637.02	0.36	w3927
2894.91	2291.60	1.26	w3069	15445.74	42484.53	0.36	w3069
1162.90	1058.22	1.10	w0162	5614.02	15712.25	0.36	w0162
494.27	467.07	1.06	w0564	1311.37	3713.19	0.35	w0564
2542.41	5907.71	0.43	lar	367.19	1045.32	0.35	lar
145.60	352.49	0.41	agaB	174.76	500.90	0.35	agaB
360.63	406.91	0.89	w0356	893.83	2593.46	0.34	w0356
146.66	228.04	0.64	ptrB	583.62	1710.66	0.34	ptrB
89.69	85.82	1.05	tdcA	140.71	420.20	0.33	tdcA
1569.84	1131.81	1.39	w0005	854.54	2591.31	0.33	w0005
949.03	723.77	1.31	w2820	711.72	2231.62	0.32	w2820
382.29	200.74	1.90	racC	145.56	473.98	0.31	racC
966.21	528.97	1.83	w1323	814.12	2717.50	0.30	w1323
2804.26	3141.20	0.89	tolQ	7715.84	26048.34	0.30	tolQ
349.28	732.94	0.48	w0535	238.50	834.39	0.29	w0535
19047.83	10107.02	1.88	w2546	13321.17	46691.65	0.29	w2546
580.46	495.64	1.17	w0553	734.83	2584.37	0.28	w0553
213.09	433.84	0.49	w1426	136.41	525.28	0.26	w1426
28409.79	32349.40	0.88	glpT	25094.68	97868.64	0.26	glpT
271.65	813.15	0.33	sapC	354.66	1417.55	0.25	sapC
1502.52	1107.42	1.36	w2597	631.01	2727.88	0.23	w2597
274.59	176.87	1.55	ais	706.69	3093.84	0.23	ais
191.80	216.29	0.89	celA	864.54	3806.84	0.23	celA
109.10	59.92	1.82	ppdB	38.01	182.15	0.21	ppdB
249.88	204.52	1.22	agaC	80.45	386.08	0.21	agaC
56.62	13.68	4.14	hrpA	546.07	2814.78	0.19	hrpA
182.61	92.19	1.98	tdcR	62.07	330.94	0.19	tdcR
5374.56	5767.37	0.93	spoU	10944.76	60533.88	0.18	spoU
456.55	279.26	1.63	w0549	956.45	5470.42	0.17	w0549
195.25	128.74	1.52	agaW	77.80	464.51	0.17	agaW
556.82	343.19	1.62	w0548	783.67	4816.90	0.16	w0548
177.02	182.96	0.97	alpA	37.14	237.77	0.16	alpA

230.79 63.00 3.66 hybF | 85.60 616.14 0.14 hybF

Gene names written in bold letters are SOS response genes; M15: 15 min exposure to MMC; M40: 40 min exposure to MMC

Table 7. Gene expressions in DM800 and DM803 when exposed to MMC

5

Gene name	b number	DM800 MMC	control	ratio	DM803 MMC	DM803 control	ratio
recN		98536.3	11895.5	8.3	2454.6	2089.9	1.2
hslS		11287.9	1677.7	6.7	426.0	370.9	1.1
dinI		10515.2	1564.7	6.7	780.0	516.8	1.5
sulA		13936.2	2416.8	5.8	1167.7	801.5	1.5
w1816	b1848	18614.3	3231.0	5.8	124.7	900.3	0.1
hslT		21578.6	5358.3	4.0	1486.7	962.6	1.5
lexA		22768.8	6375.1	3.6	2935.3	2950.2	1.0
w0522	b0531	8750.6	2628.4	3.3	1299.3	752.0	1.7
w1815	b1847	16206.3	4874.5	3.3	236.0	660.6	0.4
w2141	b2181	2156.1	691.5	3.1	0.0	934.4	0.0
recA		34375.8	11386.6	3.0	8505.3	6677.0	1.3
smgA		1614.6	558.5	2.9	591.3	624.8	0.9
cspA		12566.2	4347.9	2.9	52966.2	42356.6	1.3
ecpD		227.5	80.4	2.8	384.4	287.9	1.3
w3019	b3080	802.6	288.9	2.8	970.7	621.6	1.6
w2878	b2939	1151.8	439.8	2.6	0.0	373.7	0.0
entD		491.1	189.4	2.6	491.6	328.8	1.5
fhuC		818.1	321.7	2.5	967.4	712.6	1.4
w1201	b1228	4623.0	1874.2	2.5	2598.5	2567.4	1.0
w2999	b3059	1710.6	709.1	2.4	1279.4	1182.6	1.1
caiB		326.5	139.5	2.3	834.2	806.6	1.0
infA		8186.4	3504.9	2.3	2964.1	3046.5	1.0
uvrA		33530.3	14452.9	2.3	6731.4	5941.1	1.1
w2879	b2940	3946.5	1727.3	2.3	281.8	1248.5	0.2
insB_2		6522.2	2894.3	2.3	4566.2	4307.9	1.1
dinD		5316.7	2385.6	2.2	3182.2	2522.4	1.3
secG		6754.2	3076.6	2.2	9867.2	8993.5	1.1
priC		341.8	156.1	2.2	1151.2	509.8	2.3
w0561	b0571	11989.7	5479.6	2.2	0.0	636.5	0.0
exbD		7165.4	3289.3	2.2	1867.7	1621.7	1.2
umuC		5697.9	2659.8	2.1	1014.3	908.3	1.1
mioC		8113.1	3804.3	2.1	3218.1	2444.7	1.3
insB_1		6065.0	2854.2	2.1	4533.2	4432.3	1.0
trkH		17795.3	8430.8	2.1	2293.2	1959.1	1.2
w1345	b1374	1026.3	487.6	2.1	0.0	522.9	0.0
dniR		1878.7	900.0	2.1	1427.1	899.9	1.6
uvrB		14508.7	6960.8	2.1	5425.4	5179.7	1.0
insA_4		2298.2	1111.8	2.1	939.3	1747.8	0.5
ruvA		10492.5	5134.4	2.0	1819.8	1918.2	0.9
appY		815.1	400.3	2.0	593.1	340.6	1.7
xseA		13206.6	6494.6	2.0	17733.8	8677.0	2.0
w0224	b0231	6686.9	3310.7	2.0	2353.4	1760.8	1.3
w3139	b3199	11174.8	5555.3	2.0	2171.5	3228.1	0.7
w2512	b2559	24511.0	12207.1	2.0	1537.4	1511.4	1.0
entF		179.8	91.5	2.0	494.1	339.5	1.5
glnK		408.6	208.4	2.0	571.9	212.8	2.7
insB_4		6575.4	3364.4	2.0	2823.1	2738.2	1.0

mpA		5250.2	2686.3	2.0	1722.3	2000.1	0.9
pheP		2467.2	1263.6	2.0	2974.0	2187.8	1.4
w0491	b0500	508.2	925.1	0.5	176.3	96.9	1.8
w4088	b4183	384.5	701.1	0.5	777.9	583.0	1.3
w0570	b0580	1956.2	3578.9	0.5	867.8	197.6	4.4
w0221	b0228	1612.4	2950.5	0.5	326.2	420.6	0.8
w1347	b1376	5383.3	9852.6	0.5	1194.3	1736.1	0.7
rem		540.5	989.7	0.5	0.0	0.0	
xyIF		2117.7	3885.5	0.5	4289.6	3178.7	1.3
w2284	b2325	452.5	830.6	0.5	894.8	911.0	1.0
w0627	b0637	1984.5	3647.0	0.5	0.0	299.3	0.0
nlp		487.4	895.9	0.5	1004.9	549.6	1.8
w2940	b3001	3777.0	6949.2	0.5	2725.9	1950.7	1.4
w0591	b0601	1658.1	3060.6	0.5	602.0	402.6	1.5
w0270	b0278	1175.1	2170.8	0.5	493.9	428.3	1.2
w2790	b2849	455.2	841.4	0.5	489.3	455.5	1.1
pyrI		1174.7	2171.6	0.5	752.3	509.1	1.5
w1018	b1045	2359.6	4364.5	0.5	525.2	272.8	1.9
w2709	b2767	1312.0	2440.9	0.5	597.4	383.4	1.6
w2329	b2371	594.8	1107.2	0.5	861.9	690.9	1.2
aidB		1218.9	2269.2	0.5	4969.4	3102.9	1.6
w2791	b2850	773.2	1441.3	0.5	508.9	523.1	1.0
w2605	b2659	6211.7	11600.3	0.5	3614.7	2885.4	1.3
cmtB		141.2	263.8	0.5	275.0	164.4	1.7
w3890	b3975	824.3	1541.5	0.5	607.0	403.6	1.5
gntV		1242.4	2324.3	0.5	311.4	446.1	0.7
aspA		92135.1	173324.7	0.5	10420.0	16655.6	0.6
w3210	b3268	479.3	903.6	0.5	551.4	372.6	1.5
w0271	b0279	792.9	1496.9	0.5	471.2	426.9	1.1
w1003	b1029	2145.7	4051.8	0.5	2.6	0.0	
w1017	b1044	1431.3	2703.5	0.5	1035.5	413.5	2.5
feoA		470.5	888.8	0.5	1313.4	1090.4	1.2
chpS		1657.3	3141.1	0.5	568.1	558.4	1.0
w0619	b0629	2054.3	3896.8	0.5	289.0	43.4	6.7
yjiM	b4357	1966.9	3732.7	0.5	2090.3	1457.0	1.4
w2816	b2876	1687.7	3205.6	0.5	1046.1	858.8	1.2
w3272	b3337	587.3	1115.5	0.5	430.3	359.9	1.2
w3443	b3507	446.9	849.1	0.5	591.4	598.3	1.0
w0351	b0359	1435.0	2731.1	0.5	0.0	72.1	0.0
w1527	b1556	253.0	482.4	0.5	726.3	421.9	1.7
w0263	b0271	955.9	1824.5	0.5	484.0	330.6	1.5
csgA		522.9	998.6	0.5	703.7	599.6	1.2
fimZ		536.1	1024.6	0.5	424.3	343.7	1.2
w2314	b2356	581.8	1113.3	0.5	1114.7	731.5	1.5
w2596	b2649	370.7	710.0	0.5	1024.4	605.2	1.7
yigJ	b3823	542.9	1039.7	0.5	1372.3	817.5	1.7
w1025	b1052	1003.1	1922.7	0.5	177.0	90.5	2.0
w1359	b1388	2953.5	5661.9	0.5	328.5	262.2	1.3
w0673	b0690	1558.5	2992.7	0.5	249.4	313.2	0.8
fixX		313.0	601.3	0.5	1259.9	891.9	1.4
w4093	b4188	953.2	1839.0	0.5	901.6	606.7	1.5
w0674	b0691	1466.3	2840.2	0.5	129.0	350.8	0.4
w2400	b2447	1214.0	2355.8	0.5	810.8	406.3	2.0
w1963	b2004	537.0	1043.7	0.5	667.7	598.5	1.1
w3974	b4066	455.1	884.6	0.5	816.0	462.9	1.8
w1242	b1271	3047.5	5958.6	0.5	1870.6	336.3	5.6
w0525	b0534	930.8	1822.0	0.5	829.3	490.0	1.7

w4132	b4227	1602.0	3151.2	0.5	3125.1	2509.8	1.2
w3304	b3369	428.9	850.8	0.5	728.3	701.9	1.0
w2228	b2269	272.0	540.0	0.5	482.1	399.3	1.2
chaB		721.8	1433.4	0.5	177.0	0.0	
cydA		699.9	1392.5	0.5	963.9	305.9	3.2
melR		5082.0	10140.2	0.5	1511.6	1528.2	1.0
w1004	b1030	1265.7	2528.7	0.5	27.8	163.0	0.2
hofG		315.2	630.0	0.5	694.2	767.7	0.9
w1429	b1458	360.8	721.2	0.5	711.8	488.3	1.5
w0299	b0307	9464.0	18943.1	0.5	2048.3	2846.3	0.7
rpiR		748.7	1502.6	0.5	536.3	419.6	1.3
celD		2009.3	4041.7	0.5	1094.3	303.1	3.6
w0801	b0825	2801.7	5636.8	0.5	920.2	417.5	2.2
w0241	b0249	712.2	1434.3	0.5	772.1	649.4	1.2
w0621	b0631	1293.5	2605.4	0.5	0.0	170.6	0.0
putP		12834.1	25859.7	0.5	5693.7	6221.7	0.9
w4099	b4194	377.9	764.4	0.5	709.5	539.0	1.3
prsA		535.4	1091.6	0.5	24.5	0.0	
hybD		177.8	363.8	0.5	510.9	573.9	0.9
sapB		501.3	1027.1	0.5	258.8	0.0	
w3821	b3901	412.4	845.8	0.5	945.9	539.7	1.8
w1459	b1488	417.7	857.7	0.5	994.9	740.6	1.3
agaD		222.7	459.0	0.5	966.6	465.2	2.1
ccmD		646.1	1332.0	0.5	92.3	64.5	1.4
cpsG		1070.1	2212.1	0.5	0.0	8.9	0.0
relB		1002.3	2072.5	0.5	0.0	92.3	0.0
w2809	b2869	3556.9	7365.3	0.5	1264.0	1453.8	0.9
w0824	b0848	1242.8	2590.0	0.5	370.9	350.1	1.1
osmE		2901.6	6072.9	0.5	350.0	900.3	0.4
w0362	b0370	1234.6	2593.0	0.5	0.0	0.0	
w1927	b1963	481.3	1014.0	0.5	824.4	612.6	1.3
w0211	b0218	639.9	1353.7	0.5	198.7	321.6	0.6
w0237	b0245	855.9	1815.7	0.5	355.4	461.1	0.8
w2592	b2645	586.0	1247.1	0.5	1032.5	566.6	1.8
phnH		987.7	2147.0	0.5	1103.3	937.7	1.2
flgA		577.4	1255.7	0.5	1482.3	1396.3	1.1
w2595	b2648	377.1	821.4	0.5	647.4	431.6	1.5
w2600	b2654	792.5	1732.9	0.5	222.5	309.6	0.7
pheL		93.3	205.1	0.5	256.4	258.9	1.0
w3049	b3107	1146.6	2524.7	0.5	53.4	180.9	0.3
w1031	b1058	1045.4	2305.9	0.5	188.9	101.9	1.9
w0295	b0303	473.5	1045.0	0.5	468.2	461.2	1.0
marB		872.9	1928.8	0.5	843.0	130.7	6.5
w0665	b0682	786.7	1744.6	0.5	0.0	592.3	0.0
w1016	b1043	1399.6	3109.1	0.5	419.6	203.5	2.1
w0298	b0306	8569.6	19066.7	0.4	1722.6	2401.5	0.7
w0812	b0836	1381.4	3073.9	0.4	356.8	314.2	1.1
w2026	b2067	1456.5	3270.0	0.4	22.0	337.9	0.1
w0715	b0732	1050.7	2361.6	0.4	154.8	30.1	5.1
pyrL		689.6	1552.6	0.4	520.5	338.5	1.5
menE		1802.6	4058.7	0.4	4640.7	928.2	5.0
mb		703.1	1594.1	0.4	399.4	0.0	
fucR		1770.9	4026.5	0.4	2264.4	1572.6	1.4
w2818	b2878	278.7	634.2	0.4	787.6	511.9	1.5
acpD		1990.2	4539.4	0.4	0.0	287.0	0.0
w0489	b0498	332.5	758.7	0.4	137.3	183.4	0.7
ppdA		177.3	405.0	0.4	662.6	505.3	1.3
w1966	b2007	488.9	1118.3	0.4	1163.2	680.8	1.7

"no template"		333.4	764.2	0.4	470.2	368.3	1.3
w2401	b2448	1085.4	2497.9	0.4	1160.0	715.4	1.6
w4094	b4189	1505.9	3477.9	0.4	1022.3	932.0	1.1
dicC		543.4	1266.3	0.4	319.5	67.2	4.8
w0286	b0294	844.0	1986.6	0.4	498.7	447.4	1.1
selB		341.9	808.9	0.4	7781.6	7090.5	1.1
w2733	b2789	3589.0	8496.9	0.4	5638.1	3562.2	1.6
"no template"		301.6	716.0	0.4	408.2	329.1	1.2
trpC		10007.8	23940.6	0.4	421.9	2562.2	0.2
relF		449.8	1077.5	0.4	0.0	0.0	
w1318	b1347	1107.0	2655.3	0.4	1.4	308.5	0.0
agaV		202.3	490.1	0.4	673.0	669.2	1.0
w1002	b1028	1315.8	3189.3	0.4	288.3	62.7	4.6
w0685	b0702	1124.7	2728.6	0.4	198.6	229.5	0.9
potH		500.1	1228.0	0.4	641.9	243.8	2.6
w2399	b2446	853.3	2107.1	0.4	770.2	485.4	1.6
metA		991.4	2449.4	0.4	2026.6	1816.4	1.1
lytB		311.0	770.5	0.4	353.9	181.5	1.9
w2987	b3047	638.0	1593.1	0.4	509.8	303.9	1.7
w0552	b0562	1225.7	3084.4	0.4	0.0	588.7	0.0
w1846	b1878	382.4	968.1	0.4	1322.3	1154.7	1.1
dicB		533.3	1351.9	0.4	260.1	54.8	4.7
w1005	b1031	970.4	2465.9	0.4	0.0	106.0	0.0
w2587	b2640	509.5	1301.7	0.4	557.1	371.8	1.5
w1260	b1289	963.5	2507.3	0.4	0.0	353.6	0.0
w3068	b3127	12241.5	31901.4	0.4	2383.9	3185.5	0.7
w0551	b0561	2143.7	5619.4	0.4	477.1	864.7	0.6
w2599	b2653	616.9	1633.9	0.4	456.5	107.8	4.2
w0569	b0579	912.6	2444.1	0.4	673.5	612.3	1.1
fruR		7146.3	19274.4	0.4	2076.3	2369.5	0.9
w3927	b4020	1323.8	3637.0	0.4	2685.9	2131.5	1.3
w3069	b3128	15445.7	42484.5	0.4	499.8	1153.3	0.4
w0162	b0162	5614.0	15712.2	0.4	4564.1	3889.2	1.2
w0564	b0574	1311.4	3713.2	0.4	584.5	175.8	3.3
lar		367.2	1045.3	0.4	82.1	122.0	0.7
agaB		174.8	500.9	0.3	371.8	504.1	0.7
w0356	b0364	893.8	2593.5	0.3	10.4	32.6	0.3
ptrB		583.6	1710.7	0.3	246.4	69.2	3.6
tdcA		140.7	420.2	0.3	612.3	343.4	1.8
w0005	b0005	854.5	2591.3	0.3	442.3	388.2	1.1
w2820	b2880	711.7	2231.6	0.3	904.4	609.4	1.5
racC		145.6	474.0	0.3	167.4	26.9	6.2
w1323	b1352	814.1	2717.5	0.3	297.1	235.6	1.3
tolQ		7715.8	26048.3	0.3	1213.4	1001.4	1.2
w0535	b0545	238.5	834.4	0.3	114.4	162.5	0.7
w2546	b2597	13321.2	46691.6	0.3	2567.6	4164.9	0.6
w0553	b0563	734.8	2584.4	0.3	0.0	277.1	0.0
w1426	b1455	136.4	525.3	0.3	672.5	358.6	1.9
glpT		25094.7	97868.6	0.3	7769.7	14648.6	0.5
sapC		354.7	1417.6	0.3	546.0	0.0	
w2597	b2650	631.0	2727.9	0.2	1000.0	566.4	1.8
ais		706.7	3093.8	0.2	352.9	114.6	3.1
celA		864.5	3806.8	0.2	0.0	512.6	0.0
ppdB		38.0	182.1	0.2	399.9	345.8	1.2
agaC		80.5	386.1	0.2	598.1	602.8	1.0
hrpA		546.1	2814.8	0.2	0.0	147.3	0.0

tdcR		62.1	330.9	0.2	350.0	232.7	1.5
spoU		10944.8	60533.9	0.2	4024.5	3163.9	1.3
w0549	b0559	956.4	5470.4	0.2	115.5	287.2	0.4
agaW		77.8	464.5	0.2	479.0	779.3	0.6
w0548	b0558	783.7	4816.9	0.2	327.8	85.0	3.9
alpA		37.1	237.8	0.2	366.3	269.2	1.4
hybF		85.6	616.1	0.1	332.8	456.6	0.7

Table 8. Gene expressions in DM800 and DM803 when exposed to MMC

<u>Gene name</u>	<u>b #</u>	<u>DM800- MMC</u>	<u>DM800- control</u>	<u>DM800 ratio (MMC/control)</u>	<u>DM803- MMC</u>	<u>DM803- control</u>	<u>DM803 ratio (MMC/control)</u>
							<u>l)</u>
chaB		721.8	1433.4	0.5	177.0	0.0	
gusC		1069.6	1178.1	0.9	39.2	0.0	
prsA		535.4	1091.6	0.5	24.5	0.0	
rnb		703.1	1594.1	0.4	399.4	0.0	
rspB		987.2	1520.0	0.6	480.4	0.0	
sapB		501.3	1027.1	0.5	258.8	0.0	
sapC		354.7	1417.6	0.3	546.0	0.0	
uxaB		2051.4	2423.2	0.8	527.4	0.0	
w0367	b0375	1010.4	1827.8	0.6	179.0	0.0	
w0492	b0501	717.2	769.2	0.9	85.3	0.0	
w0521	b0530	425.3	746.2	0.6	169.9	0.0	
w0537	b0547	711.1	851.5	0.8	87.9	0.0	
w0544	b0554	602.7	558.1	1.1	98.8	0.0	
w1114	b1141	1523.8	1839.4	0.8	30.1	0.0	
w0508	b0517	2160.9	2528.0	0.9	480.7	1.0	457.9
w0541	b0551	542.2	591.7	0.9	140.0	7.6	18.4
molR		4939.4	4515.9	1.1	5518.5	513.6	10.7
w2514	b2561	7715.4	7319.4	1.1	23955.8	2367.6	10.1
molR		3903.5	3852.3	1.0	2355.4	243.6	9.7
w1329	b1358	1909.4	1872.1	1.0	649.0	75.2	8.6
w1381	b1410	2104.8	1895.7	1.1	2603.5	309.6	8.4
w1088	b1115	3071.3	4218.6	0.7	1543.9	184.8	8.4
dsdX		2030.3	2282.9	0.9	1833.5	227.0	8.1
thrS		277.1	318.4	0.9	1498.9	196.8	7.6
w0873	b0898	2210.6	1636.1	1.4	1001.4	132.9	7.5
w0617	b0627	4834.3	7145.9	0.7	2171.5	301.1	7.2
w3258	b3323	4107.7	3439.8	1.2	9485.2	1318.6	7.2
w0999	b1025	1233.1	888.5	1.4	199.5	28.3	7.1
rspA		1862.1	2002.0	0.9	1297.2	191.4	6.8
w0843	b0867	2593.1	2957.8	0.9	1333.5	197.9	6.7
w0619	b0629	2054.3	3896.8	0.5	289.0	43.4	6.7
marB		872.9	1928.8	0.5	843.0	130.7	6.5
w1128	b1155	865.4	1384.1	0.6	157.3	24.8	6.3
napH		1548.4	1771.5	0.9	2445.8	392.2	6.2
racC		145.6	474.0	0.3	167.4	26.9	6.2
w1266	b1295	1834.3	3178.5	0.6	1111.8	178.6	6.2
w2857	b2918	2013.5	2025.3	1.0	2905.4	475.0	6.1
pfkB		5266.9	4949.8	1.1	1148.4	188.9	6.1
w1309	b1338	3612.6	4271.9	0.8	3943.8	660.9	6.0
napG		4522.3	4522.5	1.0	5243.5	900.1	5.8
w1242	b1271	3047.5	5958.6	0.5	1870.6	336.3	5.6
w0991	b1017	2292.7	1982.0	1.2	564.2	101.9	5.5
vsr		5927.8	3617.9	1.6	3758.3	687.4	5.5

w0996	b1022	954.0	881.7	1.1	406.4	74.4	5.5
w0957	b0983	1903.7	1772.1	1.1	1280.4	234.3	5.5
w1287	b1316	4404.7	4724.2	0.9	481.3	89.0	5.4
w1113	b1140	2063.9	2466.4	0.8	1674.5	315.1	5.3
w1168	b1195	1068.7	1662.1	0.6	105.1	19.9	5.3
w0918	b0943	964.3	1122.4	0.9	166.1	31.9	5.2
hslJ		1701.0	1353.6	1.3	952.9	183.4	5.2
fdnH		2916.7	3353.8	0.9	1551.5	299.3	5.2
w0715	b0732	1050.7	2361.6	0.4	154.8	30.1	5.1
menE		1802.6	4058.7	0.4	4640.7	928.2	5.0
w0511	b0520	2664.2	3109.8	0.9	2221.0	452.5	4.9
w2508	b2555	3575.2	2467.8	1.4	5237.2	1078.9	4.9
pspC		670.6	809.3	0.8	370.0	76.5	4.8
w0767	b0791	2562.9	3359.1	0.8	2376.4	497.9	4.8
dicC		543.4	1266.3	0.4	319.5	67.2	4.8
dicB		533.3	1351.9	0.4	260.1	54.8	4.7
chaC		3102.1	3008.4	1.0	1480.1	314.6	4.7
w0387	b0395	2772.4	3361.1	0.8	1676.5	362.0	4.6
w0986	b1012	1670.0	1494.6	1.1	1627.0	352.8	4.6
w1002	b1028	1315.8	3189.3	0.4	288.3	62.7	4.6
w0570	b0580	1956.2	3578.9	0.5	867.8	197.6	4.4
w1972	b2013	2191.0	2234.9	1.0	1878.6	435.2	4.3
gatR		16844.2	12487.4	1.3	2559.5	598.3	4.3
w2582	b2634	3769.3	4821.9	0.8	3379.2	793.9	4.3
w2599	b2653	616.9	1633.9	0.4	456.5	107.8	4.2
w0793	b0817	9409.7	8201.4	1.1	8721.9	2060.6	4.2
w3260	b3325	1313.1	1350.2	1.0	544.2	131.4	4.1
w0888	b0913	3553.0	2732.1	1.3	3682.2	899.6	4.1
w0446	b0454	1294.0	1873.1	0.7	1977.3	483.5	4.1
xthA		7322.0	7765.8	0.9	3014.3	750.9	4.0
w2574	b2626	1092.4	1842.3	0.6	1984.7	494.8	4.0
w1211	b1240	2015.0	1589.9	1.3	685.0	171.1	4.0
btuC		5381.6	3197.6	1.7	3189.6	804.9	4.0
osmB		1731.8	2164.1	0.8	1092.6	281.1	3.9
w1308	b1337	1961.3	2585.3	0.8	2106.4	544.5	3.9
w1210	b1239	1397.2	1027.0	1.4	1996.6	516.3	3.9
w0616	b0626	2728.7	3046.4	0.9	840.4	217.6	3.9
w0548	b0558	783.7	4816.9	0.2	327.8	85.0	3.9
dsrB		803.9	1220.8	0.7	1792.6	471.9	3.8
w1382	b1411	2709.0	2309.7	1.2	2107.7	554.9	3.8
alkB		1424.0	1719.3	0.8	437.2	115.3	3.8
pyrF		4511.4	3724.1	1.2	3376.5	903.9	3.7
w1292	b1321	12154.3	8766.6	1.4	1109.1	305.2	3.6
w1331	b1360	2085.1	2530.6	0.8	1174.9	324.1	3.6
w0764	b0788	2579.2	2813.6	0.9	2474.3	682.9	3.6
mcrA		2912.9	3609.3	0.8	476.8	132.0	3.6
celD		2009.3	4041.7	0.5	1094.3	303.1	3.6
w3213	b3271	1521.2	1151.6	1.3	2042.2	570.1	3.6
w0998	b1024	2410.5	1957.8	1.2	2866.7	803.1	3.6
ptrB		583.6	1710.7	0.3	246.4	69.2	3.6
malY		3296.2	3398.9	1.0	876.6	250.7	3.5
prfH		632.2	648.7	1.0	212.1	61.6	3.4
w3013	b3074	2711.4	2653.2	1.0	2576.4	760.8	3.4
w3259	b3324	840.9	1191.9	0.7	942.0	278.2	3.4
w0719	b0736	2884.4	2820.3	1.0	866.1	257.5	3.4
w0765	b0789	3699.1	3911.8	0.9	2615.0	783.7	3.3
w0564	b0574	1311.4	3713.2	0.4	584.5	175.8	3.3
recT		1612.4	1717.9	0.9	303.3	91.9	3.3

w0923	b0948	9321.0	6900.1	1.4	4604.2	1402.6	3.3
w0817	b0841	3320.7	3960.8	0.8	3196.3	975.8	3.3
w3086	b3145	1310.1	1251.5	1.0	1040.6	317.9	3.3
bisZ		6146.6	4998.7	1.2	4600.8	1407.5	3.3
thrL		1057.1	1045.9	1.0	1851.9	566.6	3.3
w0497	b0506	9030.8	9211.8	1.0	5078.5	1555.4	3.3
w1971	b2012	1758.9	1667.2	1.1	691.8	213.9	3.2
w0540	b0550	2028.9	1744.8	1.2	1549.1	480.5	3.2
w1630	b1660	5454.2	5556.5	1.0	4639.8	1450.4	3.2
cpsB		3575.1	4298.5	0.8	1022.1	322.1	3.2
cydA		699.9	1392.5	0.5	963.9	305.9	3.2
hisQ		1785.2	2681.4	0.7	608.3	194.5	3.1
hisC		6681.3	9140.8	0.7	846.1	270.9	3.1
w0500	b0509	1305.4	1278.1	1.0	419.4	134.9	3.1
w1174	b1201	4364.8	3477.3	1.3	4444.2	1441.8	3.1
ais		706.7	3093.8	0.2	352.9	114.6	3.1
w0779	b0803	1650.0	1698.6	1.0	333.4	108.6	3.1
w1239	b1268	2079.2	1408.0	1.5	3135.3	1028.5	3.0
purB		18600.8	19192.0	1.0	5778.2	1902.4	3.0
pth		3170.2	2879.8	1.1	961.3	316.5	3.0
w0650	b0660	22565.4	15873.4	1.4	6943.2	2288.0	3.0
cheW		1753.3	1660.5	1.1	16269.0	5397.0	3.0
w1399	b1428	3396.4	3241.9	1.0	1677.5	560.3	3.0
w0961	b0987	879.4	968.1	0.9	432.4	144.6	3.0
w0936	b0961	1274.0	1621.9	0.8	613.8	205.8	3.0
w0502	b0511	5155.5	5061.0	1.0	2862.5	964.1	3.0
w0822	b0846	1673.8	2457.6	0.7	600.4	202.4	3.0
entC		152.5	97.0	1.6	611.3	206.3	3.0
sdhC		844.1	1314.1	0.6	1633.1	553.6	2.9
w0509	b0518	3075.7	3295.1	0.9	1828.1	620.4	2.9
xasA		1931.3	2558.8	0.8	1157.5	394.7	2.9
hyaE		676.4	1025.4	0.7	755.3	258.0	2.9
w0539	b0549	747.5	889.9	0.8	439.3	151.8	2.9
w1115	b1142	1661.8	2188.1	0.8	192.4	66.7	2.9
hycH		288.2	502.4	0.6	665.1	232.8	2.9
glnB		1596.2	2017.4	0.8	7231.5	2542.3	2.8
w1302	b1331	99632.3	107376.2	0.9	41005.5	14416.5	2.8
cysA		2279.2	2408.4	0.9	1175.7	414.3	2.8
codB		4134.2	2988.3	1.4	16969.3	6059.8	2.8
rbsD		765.7	795.2	1.0	759.5	271.3	2.8
w3180	b3240	1863.2	1194.4	1.6	582.5	208.5	2.8
gefL		560.6	561.3	1.0	702.5	254.1	2.8
w0958	b0984	2457.8	4067.5	0.6	1021.8	369.7	2.8
aroC		11775.3	9917.9	1.2	10669.1	3864.7	2.8
motA		2939.6	2956.1	1.0	11679.2	4245.3	2.8
w3025	b3086	1433.6	1291.0	1.1	1758.4	650.5	2.7
hemK		1309.8	2201.0	0.6	710.0	263.6	2.7
glnK		408.6	208.4	2.0	571.9	212.8	2.7
molR		2331.5	2500.4	0.9	1737.4	647.3	2.7
w1130	b1157	1347.6	1890.5	0.7	475.1	179.1	2.7
w0128	b0128	1773.0	1996.8	0.9	7243.6	2738.9	2.6
potH		500.1	1228.0	0.4	641.9	243.8	2.6
w0649	b0659	8552.5	6250.8	1.4	2846.7	1095.1	2.6
cspB		1529.3	1353.4	1.1	4972.3	1916.6	2.6
w1030	b1057	1588.1	2509.8	0.6	267.4	103.5	2.6
purU		7223.2	5962.4	1.2	5262.9	2036.9	2.6
pspD		4123.9	3232.2	1.3	3232.5	1251.9	2.6
w1367	b1396	1421.9	1262.8	1.1	402.2	155.9	2.6

vacJ		3901.7	3263.7	1.2	3362.3	1305.4	2.6
rimL		1674.8	2976.5	0.6	322.5	125.2	2.6
w2436	b2483	1505.8	2310.6	0.7	1501.8	584.0	2.6
w1969	b2010	3303.8	2815.0	1.2	3061.5	1190.7	2.6
w1499	b1528	2954.5	3022.6	1.0	1198.3	471.0	2.5
w0608	b0618	2142.0	2560.7	0.8	1265.6	499.3	2.5
acrB		300.1	343.2	0.9	702.9	277.8	2.5
fadL		11366.6	11834.8	1.0	6211.9	2462.1	2.5
deoR		544.1	807.2	0.7	1026.9	408.3	2.5
ccmF		4594.5	4971.3	0.9	3393.8	1352.8	2.5
w0473	b0482	1714.2	2540.3	0.7	2861.6	1141.4	2.5
w1017	b1044	1431.3	2703.5	0.5	1035.5	413.5	2.5
sdiA		624.8	836.4	0.7	280.1	112.0	2.5
w0845	b0869	3115.9	3364.6	0.9	2606.5	1044.4	2.5
w0702	b0719	1662.1	2274.0	0.7	163.0	65.5	2.5
nohA		6246.0	6739.5	0.9	3562.3	1439.6	2.5
w0652	b0662	3324.9	3936.8	0.8	2463.7	999.4	2.5
w0538	b0548	667.7	876.1	0.8	242.3	98.9	2.4
w2124	b2164	2328.3	2341.3	1.0	795.8	324.9	2.4
w1336	b1365	976.6	1330.0	0.7	832.5	340.1	2.4
w0507	b0516	2900.6	3116.9	0.9	1473.0	602.3	2.4
ccmB		1241.9	1597.6	0.8	518.1	211.9	2.4
marA		3033.3	3218.3	0.9	1832.6	751.7	2.4
dcp		17199.9	15990.1	1.1	9211.8	3786.9	2.4
dacB		385.1	673.3	0.6	399.3	164.3	2.4
w0794	b0818	6395.1	5722.4	1.1	4159.6	1718.0	2.4
w0692	b0709	7727.5	8615.3	0.9	1950.3	814.0	2.4
trpL		646.3	734.9	0.9	93.1	39.0	2.4
w1284	b1313	3976.7	4821.7	0.8	282.3	118.5	2.4
uraA		10187.2	8537.2	1.2	9902.4	4158.1	2.4
w0980	b1006	2643.2	2348.5	1.1	1308.5	552.5	2.4
w3017	b3078	9857.5	6296.2	1.6	6341.6	2689.7	2.4
w1353	b1382	928.5	824.8	1.1	370.0	157.8	2.3
sdaA		13413.9	11886.2	1.1	4508.6	1928.7	2.3
w0635	b0645	1636.0	2598.8	0.6	957.8	409.8	2.3
rpsV		1839.1	2871.6	0.6	393.4	168.6	2.3
nuoM		43613.3	39960.8	1.1	10711.5	4591.6	2.3
w1255	b1284	3075.9	5314.3	0.6	1188.4	509.6	2.3
fdnG		9178.0	9437.7	1.0	4206.3	1807.4	2.3
mepA		3493.6	3401.6	1.0	1838.9	790.6	2.3
w2751	b2807	1852.2	1235.3	1.5	863.6	372.1	2.3
w2593	b2646	735.3	900.8	0.8	1259.4	546.5	2.3
w0249	b0257	1814.6	2321.4	0.8	918.5	399.2	2.3
w0636	b0646	4069.4	4708.4	0.9	4894.1	2129.4	2.3
w0605	b0615	3429.0	4612.5	0.7	1239.6	539.4	2.3
w1405	b1434	2637.5	2190.6	1.2	908.0	396.3	2.3
w2056	b2097	836.8	1201.3	0.7	894.6	392.8	2.3
w1175	b1202	5780.1	5190.3	1.1	3007.7	1325.5	2.3
priC		341.8	156.1	2.2	1151.2	509.8	2.3
cdsA		3469.5	2775.1	1.3	983.1	435.9	2.3
w2536	b2583	1891.7	2031.4	0.9	2102.6	932.6	2.3
xylE		1373.3	1927.2	0.7	7998.9	3551.8	2.3
w0135	b0135	855.6	837.8	1.0	621.5	276.4	2.2
dsdA		3589.2	3871.9	0.9	5594.6	2497.9	2.2
w0611	b0621	3552.7	4502.2	0.8	4335.0	1939.6	2.2
napC		1956.7	2181.8	0.9	1111.9	500.4	2.2
w1372	b1401	12135.8	22057.4	0.6	555.5	250.5	2.2
w0801	b0825	2801.7	5636.8	0.5	920.2	417.5	2.2

w2776	b2835	3905.8	3284.2	1.2	2517.3	1143.3	2.2
phpB		1409.6	1160.8	1.2	1214.4	553.1	2.2
frvA		1064.7	1379.4	0.8	1191.0	544.1	2.2
atoS		3671.2	4696.2	0.8	1737.9	793.9	2.2
nadC		17324.6	18335.7	0.9	10757.4	4919.0	2.2
w1333	b1362	1261.3	1693.1	0.7	121.3	55.6	2.2
ptsl		34232.4	31354.3	1.1	18044.1	8284.2	2.2
w2854	b2915	870.3	1215.4	0.7	662.7	306.0	2.2
pbpG		6944.6	6379.8	1.1	4932.9	2283.2	2.2
w3083	b3142	719.8	985.0	0.7	247.8	114.8	2.2
w1144	b1171	1358.2	1358.7	1.0	381.8	177.2	2.2
ldcC		1862.3	1945.6	1.0	1001.1	465.0	2.2
pspA		3173.9	2654.2	1.2	2159.0	1002.9	2.2
w2619	b2673	1006.0	1237.6	0.8	644.6	300.0	2.1
w0157	b0157	1346.3	1157.1	1.2	1018.0	474.0	2.1
w2942	b3003	1915.5	2314.6	0.8	13358.9	6228.9	2.1
w2130	b2170	5239.7	4874.9	1.1	3027.9	1425.0	2.1
artJ		551.8	666.4	0.8	602.1	283.4	2.1
w0784	b0808	3315.6	3019.7	1.1	2233.9	1052.1	2.1
w1992	b2033	14633.5	12234.7	1.2	3432.2	1618.3	2.1
w1086	b1113	3355.4	3376.0	1.0	1210.4	571.5	2.1
w4089	b4184	527.0	904.7	0.6	921.5	437.1	2.1
bioD		1185.3	1716.8	0.7	8262.4	3929.6	2.1
w1738	b1770	943.5	1171.6	0.8	745.3	355.1	2.1
hisH		3550.4	5116.9	0.7	467.4	223.7	2.1
w1404	b1433	3838.7	2998.2	1.3	3076.7	1473.4	2.1
hrsA		2276.9	2267.0	1.0	1635.6	784.4	2.1
entE		1452.0	1208.2	1.2	1455.2	698.8	2.1
w1244	b1273	4472.8	7128.7	0.6	1505.4	723.7	2.1
agaD		222.7	459.0	0.5	966.6	465.2	2.1
w0136	b0136	503.6	708.2	0.7	467.4	225.0	2.1
gip		2057.9	2249.6	0.9	911.3	438.8	2.1
w0935	b0960	9810.7	9012.0	1.1	5596.6	2698.4	2.1
w2098	b2138	2860.2	1994.6	1.4	852.8	411.9	2.1
dcuA		40430.1	43730.0	0.9	43935.5	21225.0	2.1
w2787	b2846	934.2	1289.9	0.7	461.0	222.7	2.1
tap		4650.7	3999.2	1.2	27236.5	13172.8	2.1
menF		5124.8	4998.4	1.0	1942.6	941.5	2.1
w1016	b1043	1399.6	3109.1	0.5	419.6	203.5	2.1
w0638	b0648	1838.6	2597.3	0.7	717.4	348.2	2.1
caiF		428.4	546.9	0.8	485.8	235.9	2.1
hypC		790.3	1039.8	0.8	1320.2	642.5	2.1
w0459	b0468	2299.1	1903.3	1.2	4101.2	1997.0	2.1
w2476	b2523	5684.6	5096.2	1.1	3252.2	1584.5	2.1
w3023	b3084	2445.4	2099.4	1.2	2283.8	1116.1	2.0
w0609	b0619	3301.9	3869.5	0.9	3035.2	1484.7	2.0
bglB		1785.8	1853.5	1.0	3096.8	1515.1	2.0
xseA		13206.6	6494.6	2.0	17733.8	8677.0	2.0
cls		10644.6	9607.6	1.1	3932.1	1928.5	2.0
w3401	b3466	889.6	1114.2	0.8	633.3	310.7	2.0
cynT		613.7	533.8	1.1	756.0	372.2	2.0
w2994	b3054	10114.3	7666.9	1.3	6222.5	3064.4	2.0
w0814	b0838	3255.0	4310.8	0.8	1521.7	751.0	2.0
glfB		410.7	661.8	0.6	1198.4	592.2	2.0
w1993	b2034	10553.5	9621.4	1.1	2342.5	1158.0	2.0
w1937	b1973	693.7	780.6	0.9	812.1	401.6	2.0
mreC		2248.8	2181.3	1.0	5387.3	2666.7	2.0
w3094	b3153	894.0	1318.6	0.7	549.0	272.4	2.0

w0342	b0350	4321.9	7657.0	0.6	2305.5	1144.3	2.0
w0795	b0819	5595.7	4122.2	1.4	4127.2	2049.1	2.0
w2702	b2760	6420.7	7051.8	0.9	4832.2	2405.3	2.0
w0919	b0944	1491.9	1676.6	0.9	539.4	268.8	2.0
w4011	b4103	1183.2	1937.3	0.6	1116.2	556.8	2.0
apaH		1181.6	616.9	1.9	2062.9	1031.2	2.0
w1140	b1167	993.4	1779.5	0.6	37.8	75.6	0.5
w3038	b3096	914.3	643.0	1.4	393.6	787.9	0.5
w0834	b0858	1262.4	1631.2	0.8	83.6	167.8	0.5
w1238	b1267	2724.1	2000.4	1.4	189.3	381.0	0.5
w0682	b0699	1482.3	2557.1	0.6	76.2	154.2	0.5
w3021	b3082	953.3	551.5	1.7	367.0	742.7	0.5
w2120	b2160	3360.7	3036.7	1.1	411.9	834.3	0.5
w2169	b2210	17678.9	16595.6	1.1	1325.6	2697.8	0.5
tonB		12234.9	9196.9	1.3	1263.6	2600.0	0.5
w3071	b3130	2847.9	3106.1	0.9	522.9	1076.3	0.5
hybG		197.8	231.5	0.9	277.1	570.6	0.5
w3057	b3115	8841.4	9026.7	1.0	1268.7	2628.1	0.5
w3140	b3200	9118.3	7619.6	1.2	2245.5	4659.8	0.5
w1001	b1027	2875.6	4647.4	0.6	263.1	551.1	0.5
ccmE		2672.6	3219.4	0.8	417.4	874.6	0.5
w3123	b3184	5627.8	4212.9	1.3	775.2	1627.3	0.5
w1227	b1256	2123.5	2318.3	0.9	431.6	909.1	0.5
fadD		5851.1	7371.8	0.8	602.2	1272.2	0.5
w1863	b1895	853.4	1054.2	0.8	72.7	153.7	0.5
w2973	b3033	3719.6	3936.6	0.9	612.9	1295.5	0.5
w1081	b1108	8020.2	9265.7	0.9	2167.2	4582.7	0.5
w2750	b2806	3964.6	2858.7	1.4	757.1	1602.1	0.5
w1893	b1928	5043.4	4943.2	1.0	729.8	1551.4	0.5
w0495	b0504	3013.9	2653.4	1.1	370.2	801.8	0.5
w3092	b3151	1889.9	1880.4	1.0	737.5	1603.5	0.5
w1181	b1208	7798.7	5758.2	1.4	869.2	1895.1	0.5
narV		2637.0	3155.3	0.8	184.4	404.4	0.5
truA		11793.0	9691.1	1.2	2917.6	6458.7	0.5
w1669	b1701	9241.3	8606.6	1.1	2036.0	4536.0	0.4
w2096	b2136	5573.1	5194.1	1.1	738.2	1652.5	0.4
w0560	b0570	5151.7	6014.9	0.9	864.4	1935.4	0.4
w0384	b0392	1206.8	1717.0	0.7	114.3	259.2	0.4
w3111	b3170	8628.0	6645.7	1.3	2692.4	6162.1	0.4
trpA		6982.3	5116.3	1.4	288.9	663.8	0.4
w0729	b0753	1992.8	2456.6	0.8	300.9	692.5	0.4
w0686	b0703	11364.2	10525.1	1.1	506.9	1168.5	0.4
glpB		3375.9	2533.8	1.3	3108.0	7165.1	0.4
w3069	b3128	15445.7	42484.5	0.4	499.8	1153.3	0.4
evgA		3622.0	3054.8	1.2	605.2	1403.5	0.4
hisA		12808.2	13416.1	1.0	1336.8	3117.3	0.4
w2146	b2186	10558.5	9654.4	1.1	1037.4	2426.5	0.4
w2535	b2582	2865.9	4526.7	0.6	393.5	921.4	0.4
pntB		20402.7	20359.0	1.0	1205.5	2840.2	0.4
w2195	b2236	1406.5	728.5	1.9	234.1	555.8	0.4
w1543	b1572	1282.8	1569.8	0.8	120.9	287.6	0.4
w0850	b0874	3017.5	2927.9	1.0	738.9	1765.9	0.4
btuE		2923.2	2641.6	1.1	263.9	631.0	0.4
w1395	b1424	1949.0	1623.8	1.2	410.0	980.9	0.4
w1356	b1385	2167.3	1587.5	1.4	96.8	234.3	0.4
w1213	b1242	503.3	619.9	0.8	151.7	369.1	0.4
w2361	b2408	763.3	752.9	1.0	120.0	292.1	0.4

w3104	b3163	18167.7	11453.8	1.6	13491.8	32888.9	0.4
w0748	b0772	11979.9	9899.5	1.2	1966.2	4794.2	0.4
w1192	b1219	3462.0	3677.7	0.9	438.1	1068.7	0.4
w3228	b3293	2810.8	1965.5	1.4	313.1	764.1	0.4
w2849	b2909	10514.5	10130.8	1.0	2114.0	5172.6	0.4
w1393	b1422	7786.4	7668.8	1.0	1190.6	2921.0	0.4
speG		14600.8	12145.6	1.2	2225.6	5463.2	0.4
w3041	b3099	5252.8	5281.6	1.0	471.9	1163.2	0.4
w3066	b3125	27223.3	36983.8	0.7	1092.1	2695.7	0.4
w2765	b2824	851.9	964.4	0.9	148.9	368.0	0.4
w1825	b1857	5399.7	4629.8	1.2	637.0	1579.2	0.4
w1281	b1310	3066.4	5129.5	0.6	143.7	356.9	0.4
w1117	b1144	1417.4	1639.5	0.9	108.6	270.0	0.4
w0549	b0559	956.4	5470.4	0.2	115.5	287.2	0.4
treA		11547.4	11404.7	1.0	1095.3	2737.4	0.4
w0778	b0802	1350.5	1308.3	1.0	131.7	329.4	0.4
w3129	b3190	2307.5	1992.7	1.2	340.2	852.2	0.4
w3218	b3283	3878.7	2425.8	1.6	1252.5	3153.4	0.4
w3053	b3111	6234.8	6408.0	1.0	607.4	1542.6	0.4
w0475	b0484	15214.1	16208.9	0.9	709.7	1802.9	0.4
w2117	b2157	3070.1	2873.2	1.1	533.7	1364.3	0.4
w1432	b1461	1846.2	2349.9	0.8	90.3	231.7	0.4
osmE		2901.6	6072.9	0.5	350.0	900.3	0.4
w1624	b1654	6959.3	7199.0	1.0	1247.1	3214.9	0.4
w0821	b0845	8372.0	8637.9	1.0	1748.5	4513.0	0.4
w0365	b0373	2918.6	3790.8	0.8	280.0	723.3	0.4
katE		4260.9	3635.5	1.2	516.6	1343.5	0.4
w0506	b0515	2698.3	2939.2	0.9	582.2	1515.2	0.4
w2354	b2398	2997.8	3805.3	0.8	381.5	996.5	0.4
sseA		11236.9	13524.4	0.8	1461.2	3819.1	0.4
w3091	b3150	4653.7	4947.8	0.9	926.2	2450.6	0.4
dadX		14179.7	15104.4	0.9	1014.8	2688.2	0.4
w1280	b1309	7116.1	7431.6	1.0	485.6	1300.2	0.4
w0909	b0934	5777.4	6344.0	0.9	191.5	512.9	0.4
w1253	b1282	4767.1	6492.3	0.7	342.5	917.5	0.4
w2977	b3037	10325.0	9553.3	1.1	2711.8	7333.0	0.4
w1876	b1908	7904.1	7700.0	1.0	1834.1	4974.1	0.4
w0674	b0691	1466.3	2840.2	0.5	129.0	350.8	0.4
w0496	b0505	860.2	954.3	0.9	107.7	293.4	0.4
w3085	b3144	8990.5	7250.8	1.2	2060.6	5627.0	0.4
w1815	b1847	16206.3	4874.5	3.3	236.0	660.6	0.4
w1506	b1535	1746.6	2351.1	0.7	504.8	1413.1	0.4
w1020	b1047	2890.7	3663.4	0.8	157.9	442.8	0.4
dsbB		5243.3	4561.9	1.1	649.9	1848.6	0.4
w1090	b1117	3645.1	3701.4	1.0	359.8	1025.0	0.4
w1880	b1915	1778.5	1658.4	1.1	148.1	422.8	0.4
w1138	b1165	1592.9	1932.4	0.8	88.0	251.7	0.3
w0345	b0353	23531.8	26792.5	0.9	2375.5	6866.5	0.3
ndk		13360.7	13564.1	1.0	1108.3	3204.9	0.3
w1154	b1181	3992.7	2962.9	1.3	336.5	995.2	0.3
w1146	b1173	1361.2	1800.5	0.8	35.4	105.1	0.3
pepT		4280.0	6805.4	0.6	394.8	1170.9	0.3
w0747	b0771	1020.8	1582.4	0.6	206.3	613.2	0.3
w3052	b3110	4449.2	5096.1	0.9	768.8	2307.4	0.3
w1465	b1494	1154.2	1682.8	0.7	37.4	112.5	0.3
w1173	b1200	3963.5	3786.2	1.0	875.3	2636.8	0.3
w1237	b1266	3678.6	3075.5	1.2	330.0	1008.8	0.3
w2480	b2527	5210.0	3260.6	1.6	296.4	907.5	0.3

cspF		769.7	724.6	1.1	1043.9	3221.9	0.3
w1344	b1373	12449.4	12132.6	1.0	631.0	1956.4	0.3
w0356	b0364	893.8	2593.5	0.3	10.4	32.6	0.3
w0445	b0453	3085.5	3417.8	0.9	208.0	664.3	0.3
w1671	b1703	10135.4	10732.8	0.9	617.4	1975.5	0.3
w0483	b0492	32526.5	25269.0	1.3	997.8	3218.2	0.3
w1101	b1128	8372.2	7496.5	1.1	1030.2	3328.9	0.3
w1121	b1148	13523.9	13946.7	1.0	994.8	3277.0	0.3
fumA		63238.6	79002.7	0.8	1142.5	3808.2	0.3
otsA		3996.5	3237.0	1.2	472.5	1597.7	0.3
w3049	b3107	1146.6	2524.7	0.5	53.4	180.9	0.3
w1736	b1768	5360.4	5223.0	1.0	270.9	918.4	0.3
w1111	b1138	2285.3	3266.6	0.7	172.2	586.3	0.3
w1078	b1105	7249.1	7241.5	1.0	336.8	1168.0	0.3
nadE		7651.4	8555.6	0.9	750.5	2616.3	0.3
w2863	b2924	4415.5	3933.5	1.1	763.3	2661.9	0.3
amiA		5181.5	4164.8	1.2	1454.0	5128.4	0.3
glpC		2150.9	1770.1	1.2	2910.6	10306.1	0.3
marR		1626.3	1757.0	0.9	147.1	523.2	0.3
w2132	b2172	4870.0	5975.5	0.8	339.6	1208.0	0.3
w1814	b1846	4710.1	4725.8	1.0	601.5	2148.6	0.3
w2781	b2840	2411.6	2180.0	1.1	323.1	1162.5	0.3
w0806	b0830	5660.3	7149.0	0.8	374.2	1356.2	0.3
sbmC		15497.2	13575.9	1.1	2005.1	7281.2	0.3
w2739	b2795	8350.0	12065.7	0.7	1466.6	5391.5	0.3
w1358	b1387	5568.3	6108.2	0.9	279.4	1034.1	0.3
w0842	b0866	1432.7	2445.9	0.6	54.2	201.1	0.3
w1282	b1311	6776.5	7862.7	0.9	292.4	1090.9	0.3
glk		6398.4	6720.8	1.0	675.1	2528.6	0.3
w2286	b2327	2122.7	1766.9	1.2	546.0	2070.8	0.3
w1286	b1315	2562.4	3691.1	0.7	103.1	391.1	0.3
w1355	b1384	3482.9	3590.2	1.0	90.5	350.2	0.3
w1270	b1299	5513.5	6834.7	0.8	355.7	1394.8	0.3
w2099	b2139	1029.7	906.0	1.1	106.5	418.0	0.3
w1407	b1436	749.6	682.5	1.1	14.0	54.9	0.3
w3155	b3215	913.5	1043.0	0.9	91.6	362.6	0.3
w2511	b2558	6964.6	7412.8	0.9	864.3	3453.5	0.3
pntA		18845.0	19495.1	1.0	697.9	2832.4	0.2
w3054	b3112	1774.6	2168.9	0.8	56.8	235.3	0.2
w0633	b0643	6699.0	11110.2	0.6	152.1	631.4	0.2
w0826	b0850	2356.4	2497.1	0.9	125.2	527.4	0.2
w1328	b1357	1405.4	1746.9	0.8	7.6	32.1	0.2
w0344	b0352	5703.2	9448.7	0.6	317.6	1350.6	0.2
narW		4827.2	5275.1	0.9	451.1	1919.6	0.2
w1119	b1146	5263.5	6210.8	0.8	130.3	564.5	0.2
w0629	b0639	4754.0	4261.2	1.1	1120.5	4889.2	0.2
w2879	b2940	3946.5	1727.3	2.3	281.8	1248.5	0.2
w2137	b2177	3760.4	3335.6	1.1	353.4	1573.1	0.2
w0983	b1009	1169.8	1186.2	1.0	66.5	296.6	0.2
w3050	b3108	7051.6	7672.6	0.9	390.7	1745.2	0.2
w1042	b1069	5901.1	5801.9	1.0	1017.7	4599.0	0.2
w1298	b1327	2105.4	2517.0	0.8	57.6	260.7	0.2
w3047	b3105	8464.9	9065.1	0.9	570.5	2624.0	0.2
w0668	b0685	4431.4	3088.6	1.4	318.4	1471.6	0.2
w0855	b0879	4397.7	3462.3	1.3	2098.5	9718.9	0.2
w0669	b0686	2371.1	2887.9	0.8	185.7	873.3	0.2
w1006	b1033	7514.8	8532.4	0.9	449.3	2183.1	0.2
hisD		8934.6	9271.5	1.0	490.5	2443.1	0.2

w0372	b0380	2303.5	2793.8	0.8	96.2	481.6	0.2
w1225	b1254	1455.4	1378.9	1.1	241.2	1218.1	0.2
w1137	b1164	1801.0	1518.1	1.2	53.4	280.3	0.2
w3039	b3097	10297.8	10899.1	0.9	470.9	2476.5	0.2
w1662	b1694	6288.5	6529.8	1.0	488.8	2613.0	0.2
w1129	b1156	912.1	1118.2	0.8	44.4	242.2	0.2
w0490	b0499	57928.7	59454.4	1.0	7786.3	43190.2	0.2
w0359	b0367	5960.8	7575.1	0.8	508.8	2903.2	0.2
w1004	b1030	1265.7	2528.7	0.5	27.8	163.0	0.2
ccmC		1392.8	1425.9	1.0	59.4	352.2	0.2
w0350	b0358	2511.7	3611.1	0.7	62.7	380.2	0.2
trpC		10007.8	23940.6	0.4	421.9	2562.2	0.2
w1299	b1328	2565.5	1837.8	1.4	67.1	430.4	0.2
w0478	b0487	2303.2	2595.4	0.9	88.7	587.3	0.2
w3098	b3157	5948.5	6214.2	1.0	1142.5	7663.5	0.1
hipB		1299.7	2047.8	0.6	25.8	185.6	0.1
w0688	b0705	2081.9	1363.9	1.5	122.4	883.1	0.1
w1816	b1848	18614.3	3231.0	5.8	124.7	900.3	0.1
menC		5525.9	8590.7	0.6	244.3	1823.6	0.1
narU		6244.4	6678.0	0.9	559.1	4236.9	0.1
cbl		1725.9	1351.0	1.3	119.7	941.6	0.1
w3055	b3113	1865.9	2074.1	0.9	35.3	278.5	0.1
w2869	b2930	15958.1	16895.0	0.9	246.6	1970.2	0.1
w0487	b0496	1988.4	2379.9	0.8	95.5	777.4	0.1
w2875	b2936	6843.6	7758.0	0.9	235.8	1928.8	0.1
w3160	b3220	1100.3	963.4	1.1	91.4	806.3	0.1
phoQ		8670.4	10289.4	0.8	163.6	1484.0	0.1
ldhA		9971.9	10016.8	1.0	78.0	788.4	0.1
w2101	b2141	1452.9	1851.1	0.8	52.3	529.2	0.1
w1402	b1431	4730.2	4071.1	1.2	69.9	717.8	0.1
uidB		6144.1	6476.0	0.9	395.6	4081.9	0.1
w0480	b0489	8009.4	7942.3	1.0	302.4	3174.6	0.1
w0626	b0636	3193.6	4188.0	0.8	141.5	1580.2	0.1
w1542	b1571	1051.8	1175.0	0.9	22.1	275.2	0.1
w1118	b1145	3879.7	3857.0	1.0	87.8	1247.7	0.1
w2026	b2067	1456.5	3270.0	0.4	22.0	337.9	0.1
w1325	b1354	1084.3	1857.7	0.6	13.8	241.2	0.1
w3062	b3120	841.5	1073.1	0.8	10.3	193.4	0.1
w0943	b0968	1175.1	1146.1	1.0	21.9	442.3	0.0
w3044	b3102	3025.5	3670.0	0.8	34.4	698.5	0.0
w1546	b1576	1343.3	1668.5	0.8	5.0	113.9	0.0
w0940	b0965	2874.6	2907.6	1.0	34.8	925.2	0.0
w2071	b2112	1583.8	2171.7	0.7	13.7	428.0	0.0
w0664	b0681	5986.0	8122.3	0.7	35.0	1167.7	0.0
w2012	b2053	3678.7	4198.0	0.9	62.8	2505.6	0.0
w0476	b0485	9577.5	11490.0	0.8	89.6	4163.9	0.0
w0479	b0488	2163.7	1845.1	1.2	10.7	1159.8	0.0
w0975	b1001	1642.8	1249.9	1.3	3.5	489.8	0.0
w1318	b1347	1107.0	2655.3	0.4	1.4	308.5	0.0
acpD		1990.2	4539.4	0.4	0.0	287.0	0.0
asr		819.7	1306.6	0.6	0.0	199.7	0.0
celA		864.5	3806.8	0.2	0.0	512.6	0.0
cirA		8601.6	7219.3	1.2	0.0	1161.0	0.0
cpsG		1070.1	2212.1	0.5	0.0	8.9	0.0
dedA		1510.1	2016.6	0.7	0.0	208.3	0.0
emrK		2298.7	2752.0	0.8	0.0	444.5	0.0
emrY		1567.1	1436.2	1.1	0.0	407.5	0.0
evgS		455.7	574.5	0.8	0.0	439.3	0.0

fumC		540.2	591.2	0.9	0.0	175.8	0.0
gadB		1951.2	2743.6	0.7	0.0	81.7	0.0
galS		6159.6	7496.6	0.8	0.0	1028.9	0.0
hrpA		546.1	2814.8	0.2	0.0	147.3	0.0
ogrK		1246.5	1941.6	0.6	0.0	118.2	0.0
pheM		221.1	390.9	0.6	0.0	73.8	0.0
potA		2332.9	3825.0	0.6	0.0	375.0	0.0
pspB		610.6	1027.9	0.6	0.0	86.1	0.0
relB		1002.3	2072.5	0.5	0.0	92.3	0.0
sieB		523.8	928.1	0.6	0.0	21.1	0.0
tynA		2320.6	3208.4	0.7	0.0	447.3	0.0
uidA		753.8	1174.9	0.6	0.0	149.9	0.0
w0351	b0359	1435.0	2731.1	0.5	0.0	72.1	0.0
w0371	b0379	1275.3	2035.9	0.6	0.0	243.3	0.0
w0477	b0486	2340.1	2797.3	0.8	0.0	1901.6	0.0
w0481	b0490	1952.8	2004.4	1.0	0.0	836.5	0.0
w0482	b0491	2892.8	2263.9	1.3	0.0	783.6	0.0
w0501	b0510	648.1	1159.1	0.6	0.0	55.3	0.0
w0529	b0539	764.4	968.4	0.8	0.0	548.7	0.0
w0532	b0542	753.5	1045.5	0.7	0.0	264.9	0.0
w0533	b0543	1016.7	1055.4	1.0	0.0	403.8	0.0
w0552	b0562	1225.7	3084.4	0.4	0.0	588.7	0.0
w0553	b0563	734.8	2584.4	0.3	0.0	277.1	0.0
w0561	b0571	11989.7	5479.6	2.2	0.0	636.5	0.0
w0621	b0631	1293.5	2605.4	0.5	0.0	170.6	0.0
w0627	b0637	1984.5	3647.0	0.5	0.0	299.3	0.0
w0656	b0671	2622.4	2405.6	1.1	0.0	624.8	0.0
w0665	b0682	786.7	1744.6	0.5	0.0	592.3	0.0
w0672	b0689	1354.0	1108.1	1.2	0.0	556.1	0.0
w0807	b0831	2452.0	3462.8	0.7	0.0	433.2	0.0
w0808	b0832	3057.1	3778.8	0.8	0.0	920.3	0.0
w0809	b0833	3158.7	3447.2	0.9	0.0	396.2	0.0
w0819	b0843	1481.2	2642.2	0.6	0.0	238.3	0.0
w0910	b0935	3537.6	4051.8	0.9	0.0	435.5	0.0
w0911	b0936	1362.2	1452.9	0.9	0.0	346.2	0.0
w0912	b0937	1344.2	1345.4	1.0	0.0	326.2	0.0
w0913	b0938	373.1	543.7	0.7	0.0	250.8	0.0
w0914	b0939	578.4	552.7	1.0	0.0	328.0	0.0
w0915	b0940	2161.6	1966.5	1.1	0.0	507.3	0.0
w0916	b0941	2523.2	2419.8	1.0	0.0	220.3	0.0
w0917	b0942	1050.8	1890.1	0.6	0.0	239.7	0.0
w0939	b0964	1052.3	1075.0	1.0	0.0	1051.3	0.0
w0944	b0969	2833.2	3213.2	0.9	0.0	480.3	0.0
w0955	b0981	3807.1	3809.9	1.0	0.0	529.4	0.0
w0956	b0982	986.7	520.9	1.9	0.0	161.0	0.0
w0973	b0999	2665.2	2466.9	1.1	0.0	705.6	0.0
w0977	b1003	3057.7	2353.2	1.3	0.0	428.1	0.0
w0990	b1016	698.0	1088.2	0.6	0.0	120.4	0.0
w0993	b1019	4411.3	4083.2	1.1	0.0	533.9	0.0
w0995	b1021	818.6	965.0	0.8	0.0	372.4	0.0
w0997	b1023	972.4	1064.8	0.9	0.0	17.8	0.0
w1005	b1031	970.4	2465.9	0.4	0.0	106.0	0.0
w1008	b1035	2819.0	3842.8	0.7	0.0	909.5	0.0
w1023	b1050	3725.1	3062.3	1.2	0.0	635.7	0.0
w1070	b1097	1352.2	1839.3	0.7	0.0	341.0	0.0
w1120	b1147	1528.0	2289.1	0.7	0.0	217.5	0.0
w1122	b1149	2784.5	2661.5	1.0	0.0	268.2	0.0
w1123	b1150	1299.2	1407.6	0.9	0.0	194.3	0.0

WO 01/29261

PCT/US00/28352

w1124	b1151	1190.2	962.1	1.2	0.0	240.8	0.0
w1125	b1152	2478.7	2830.5	0.9	0.0	378.3	0.0
w1169	b1196	736.1	1002.3	0.7	0.0	78.9	0.0
w1222	b1251	2361.8	2659.4	0.9	0.0	330.7	0.0
w1224	b1253	1462.6	1237.4	1.2	0.0	345.0	0.0
w1260	b1289	963.5	2507.3	0.4	0.0	353.6	0.0
w1267	b1296	9885.6	9682.9	1.0	0.0	628.8	0.0
w1268	b1297	6231.0	6335.9	1.0	0.0	1180.7	0.0
w1269	b1298	2180.4	2708.2	0.8	0.0	604.9	0.0
w1291	b1320	3070.2	3630.9	0.8	0.0	102.0	0.0
w1293	b1322	10797.3	10432.1	1.0	0.0	1328.7	0.0
w1317	b1346	1201.4	1869.7	0.6	0.0	2.7	0.0
w1327	b1356	7293.7	5900.3	1.2	0.0	99.2	0.0
w1332	b1361	979.7	1634.9	0.6	0.0	53.7	0.0
w1335	b1364	1031.9	1570.7	0.7	0.0	64.7	0.0
w1339	b1368	2401.5	2866.5	0.8	0.0	232.5	0.0
w1340	b1369	552.9	767.7	0.7	0.0	289.8	0.0
w1343	b1372	9415.8	9734.0	1.0	0.0	1484.3	0.0
w1345	b1374	1026.3	487.6	2.1	0.0	522.9	0.0
w1346	b1375	805.3	426.3	1.9	0.0	674.8	0.0
w1360	b1389	950.2	1644.0	0.6	0.0	254.5	0.0
w1368	b1397	1477.0	1619.0	0.9	0.0	175.4	0.0
w1390	b1419	1168.9	657.9	1.8	0.0	116.3	0.0
w1391	b1420	1225.3	1356.0	0.9	0.0	289.5	0.0
w1403	b1432	812.3	800.5	1.0	0.0	169.4	0.0
w1660	b1692	1390.9	1898.4	0.7	0.0	182.7	0.0
w1673	b1705	920.4	743.7	1.2	0.0	233.8	0.0
w1701	b1733	5606.3	5366.3	1.0	0.0	1445.5	0.0
w1733	b1765	8186.5	8316.3	1.0	0.0	1117.1	0.0
w1737	b1769	3089.3	3553.2	0.9	0.0	418.7	0.0
w2009	b2050	8039.2	9293.5	0.9	0.0	868.0	0.0
w2010	b2051	2505.5	4017.0	0.6	0.0	617.7	0.0
w2011	b2052	3552.4	5133.3	0.7	0.0	1374.5	0.0
w2078	b2119	1991.3	2900.8	0.7	0.0	261.7	0.0
w2133	b2173	848.4	657.5	1.3	0.0	474.2	0.0
w2138	b2178	5382.5	4913.7	1.1	0.0	1245.3	0.0
w2139	b2179	6107.9	5671.8	1.1	0.0	1804.5	0.0
w2140	b2180	7963.0	4829.8	1.6	0.0	815.7	0.0
w2141	b2181	2156.1	691.5	3.1	0.0	934.4	0.0
w2528	b2575	5332.1	6275.8	0.8	0.0	876.9	0.0
w2530	b2577	2228.3	2337.0	1.0	0.0	849.9	0.0
w2531	b2578	2185.4	2216.2	1.0	0.0	736.1	0.0
w2862	b2923	2498.1	1707.4	1.5	0.0	747.4	0.0
w2867	b2928	3001.1	1618.2	1.9	0.0	1005.1	0.0
w2868	b2929	2890.3	2306.0	1.3	0.0	552.7	0.0
w2878	b2939	1151.8	439.8	2.6	0.0	373.7	0.0
w2880	b2941	1480.2	1128.1	1.3	0.0	179.2	0.0
w3043	b3101	1869.2	2394.6	0.8	0.0	353.3	0.0
w3046	b3104	1186.2	1498.4	0.8	0.0	504.3	0.0
w3063	b3121	961.7	1080.5	0.9	0.0	148.6	0.0
w3084	b3143	977.8	1260.3	0.8	0.0	74.4	0.0
w3096	b3155	1052.3	847.0	1.2	0.0	714.4	0.0
w3130	b3191	5254.6	4119.2	1.3	0.0	989.0	0.0
w3133	b3194	579.9	528.6	1.1	0.0	257.1	0.0
w3265	b3330	748.4	885.6	0.8	0.0	78.5	0.0
xapR		4531.8	4954.8	0.9	0.0	1106.9	0.0
	b3399	5216.0	2817.8	1.9			

Table 9

Most highly expressed genes in *Synechocystis* sp. PCC6803 in minimal growth media
(BG11 + 5mM glucose).

5

Systematic Name	Gene	Function	Transcript copy in total mRNA (Average copy=1)
slr2051	cpcG	phycobilisome rod-core linker polypeptide CpcG	64.91
sll1580	cpcC	phycocyanin associated linker protein	22.71
slr0447	amiC	negative aliphatic amidase regulator	19.45
sll1070	tktA	transketolase	19.24
sll0018	cbbA	fructose-1, 6-bisphosphate aldolase	14.27
slr0011	rbcX	ND*	12.00
ssl0563	psaC	photosystem I subunit VII	11.31
slr1655	psaL	photosystem I subunit XI	10.91
sll0819	psaF	photosystem I subunit III	10.56
sll1867	psbA3	photosystem II D1 protein	10.43
sll1324	atpF	ATP synthase subunit b	10.37
sll1746	rpl12	50S ribosomal protein L12	10.13
sll1099	tufA	protein synthesis elongation factor Tu	9.48
slr0009	rbcL	ribulose biphosphate carboxylase large subunit	8.39
slr0012	rbcS	ribulose biphosphate carboxylase small subunit	8.14
sll1326	atpA	ATP synthase a subunit	7.72
slr1908		ND*	7.62
sll1578	cpcA	phycocyanin a subunit	7.60
slr2067	apcA	allophycocyanin a chain	7.51
slr2052		ND*	7.41
sll1184	ho	heme oxygenase	7.27
ssl3437	rps17	30S ribosomal protein S17	7.26
sll1786		hypothetical protein (ND*)	7.16
ssl0020	petF	ferredoxin	7.07
sll1812	rps5	30S ribosomal protein S5	7.04

* ND = not determined

Table 10

Most highly induced genes in *Synechocystis* sp. PCC6803 in BG11 media containing 5 mM glucose, with 20 min of UV-B treatment at $20 \mu\text{ES}^{-1}\text{m}^{-2}$ intensity.

Systematic Name	Gene	Function	Data /Control	STD
ssr2595	hliB	High light- inducible protein	22.7	4.7
slr1544		ND*	15.5	7.6
sll0528		ND*	12.1	3.9
sll1514	hsp17	small heat shock protein	9.9	3.9
slr1687	nblB	phycobilisome degradation protein NblB	8.2	1.9
sll1483		transforming growth factor induced protein	7.8	2.2
sll2012	rpoD	RNA polymerase sigma factor	6.3	2.0
ssl1633		CAB/ELIP/HLIP superfamily	6.0	1.0
ssl2542	hliA	high light-inducible protein	5.6	1.6
sll0846		ND*	4.7	0.9
slr1674		ND*	4.7	1.8
slr1604	ftsH	Chloroplast associated protease FtsH	4.6	1.9
slr0320		ND*	4.5	2.2
sll0306	rpoD	RNA polymerase sigma factor	4.4	1.0
slr0228	ftsH	cell division protein FtsH	4.3	1.7
slr1641	clpB	ClpB protein	4.3	1.1
ssr2016		ND*	4.2	2.2
sll1867	psbA3	photosystem II D1 protein	4.1	0.3

* ND = not determined

CLAIMS

What is claimed is:

1. A method for identifying gene expression changes within a bacterial species comprising:

- 5 (a) providing a comprehensive micro-array synthesized from DNA comprised in a bacterial species;
- (b) generating a first set of labeled probes from bacterial RNA, the RNA isolated from the bacterial species of step (a);
- (c) hybridizing the first set of labeled probes of step (b) to the
10 comprehensive micro-array of step (a), wherein hybridization results in a detectable signal generated from the labeled probe;
- (d) measuring the signal generated by the hybridization of the first set of labeled probe to the comprehensive micro-array of step (c);
- (e) subjecting the bacterial species of step (a) to a gene expression
15 altering condition whereby the gene expression profile of the bacterial species is altered to produce a modified bacterial species ;
- (f) generating a second set of labeled probes from bacterial RNA,
20 the RNA isolated from the modified bacterial species of step (e);
- (g) hybridizing the second set of labeled probes of step (f) to the comprehensive micro-array of step (a), wherein hybridization results in a detectable signal generated from the labeled probe;
- (h) measuring the signal generated by the hybridization of the
25 second set of labeled probes to the comprehensive micro-array of step (g); and
- (i) comparing signal generated from the first hybridization to the signal generated from the second hybridization to identify gene expression changes within a bacterial species.

30 2. A method for identifying gene expression changes within a bacterial species comprising:

- (a) providing a comprehensive micro-array synthesized from DNA comprised in a bacterial species ;
- (b) generating a first set of fluorescent cDNA from bacterial RNA,
35 the RNA isolated from the bacterial species of step (a);
- (c) hybridizing the first set of fluorescent cDNA of step (b) to the comprehensive micro-array of step (a), wherein hybridization

- results in a detectable signal generated from the fluorescent cDNA;
- (d) measuring the signal generated by the hybridization of the first set of fluorescent cDNA to the comprehensive micro-array of step (c);
- (e) subjecting the bacterial species of step (a) to a gene expression altering condition whereby the gene expression profile of the bacterial species is altered to produce a modified bacterial species;
- (f) generating a second set of fluorescent cDNA from bacterial RNA, the RNA isolated from the modified bacterial species of step (e);
- (g) hybridizing the second set of fluorescent cDNA of step (f) to the comprehensive micro-array of step (a), wherein hybridization results in a detectable signal generated from the fluorescent cDNA;
- (h) measuring the signal generated by the hybridization of the second set of fluorescent cDNA to the comprehensive micro-array of step (g); and
- (i) comparing signal generated from the first hybridization to the signal generated from the second hybridization to identify gene expression changes within a bacterial species .

3. A method according to either Claim 1 or 2 wherein the bacterial species is selected from the group consisting of enteric bacteria, *Bacillus*, *Acinetobacter*, *Streptomyces*, *Methylobacter*, *Pseudomonas*, *Rhodobacter* and *Synechocystis*

4. A method according to either Claim 1 or 2 wherein the signal generating label is selected from the group consisting of fluorescent moieties, chemiluminescent moieties, particles, enzymes, radioactive tags.

5. A method according to Claim 4 wherein the signal generating label is a fluorescent moiety and is selected from the group consisting of cy3 and cy5.

6. A method according to either Claim 1 or 2 wherein the comprehensive micro-array contains at least 75% of all open reading frames in the bacterial species.

7. A method according to Claim 6 wherein the comprehensive micro-array contains from about 2000 to about 6000 open reading frames.

8. A method according to either Claim 1 or 2 wherein the gene expression altering condition is selected from the group consisting of a condition

altering the genotype of the bacterial species, a condition altering the growth of the bacterial species, exposure to mutagens, antibiotics, UV light, gamma-rays, x-rays, phage, macrophages, organic chemicals, inorganic chemicals, environmental pollutants, heavy metals, changes in temperature, changes in pH, conditions producing oxidative damage, DNA damage, anaerobiosis, depletion or
5 addition of nutrients, addition of a growth inhibitor, and desiccation.

9. A method for identifying gene expression changes within a genome comprising:

- 10 (a) providing a comprehensive micro-array synthesized from DNA comprised in a prokaryotic or eukaryotic species;
- (b) generating a control set of fluorescent cDNA from total or polyadenylated RNA, the RNA isolated from the species of step (a), the fluorescent cDNA comprising at least one first fluorescent label and at least one different second fluorescent
15 label;
- (c) mixing the control set of fluorescent cDNA labeled with the at least one first label with the control set of fluorescent cDNA labeled with the at least second first label to form a dual labeled control cDNA;
- 20 (d) hybridizing the dual labeled control set of fluorescent cDNA of step (c) to the comprehensive micro-array of step (a), wherein hybridization results in a detectable signal generated from the fluorescent cDNA;
- (e) measuring the signal generated by the hybridization of the dual
25 labeled control set of fluorescent cDNA to the comprehensive micro-array of step (c);
- (f) subjecting the prokaryote or eukaryote of step (a) to a gene expression altering condition whereby the gene expression profile of the prokaryote or eukaryote is altered to produce a modified prokaryote or eukaryote ;
30
- (g) generating an experimental set of fluorescent cDNA from total or polyadenylated RNA, the RNA isolated from the modified prokaryote or eukaryote of step (e), the fluorescent cDNA comprising the first fluorescent label and the different second fluorescent label to step (b);
35
- (h) mixing the experimental set of fluorescent cDNA labeled with the at least one first label with the experimental set of

fluorescent cDNA labeled with the at least second first label to form a dual labeled experimental cDNA;

- 5 (i) hybridizing the experimental set of fluorescent cDNA of step (h) to the comprehensive micro-array of step (a), wherein hybridization results in a detectable signal generated from the fluorescent cDNA;
- (j) measuring the signal generated by the hybridization of the second set of fluorescent cDNA to the comprehensive micro-array of step (g); and
- 10 (k) comparing signal generated from the dual labeled control hybridization with the dual labeled experimental hybridization to identify gene expression changes within a prokaryotic or eukaryotic species.

15 10. A method according to Claim 9 wherein the first fluorescent label and the second fluorescent label is independently selected from the group consisting of cy3 and cy5.

11. A method according to Claim 9 wherein the prokaryotic or eukaryotic genome is comprised within an organism selected from the group consisting of enteric bacteria, *Bacillus*, *Acinetobacter*, *Streptomyces*, *Methylobacter*,
20 *Pseudomona*, cyanobacteria, yeasts, filamentous fungi, plant cells and animal cells.

12. A method according to Claim 11 wherein yeast are selected from the group consisting of *Saccharomyces*, *Zygosaccharomyces*, *Kluyveromyces*, *Candida*, *Hansenula*, *Debaryomyces*, *Mucor*, *Pichia* and *Torulopsis*.

25 13. A method according to Claim 11 wherein cyanobacteria are selected from the group consisting of *Rhodobacter* and *Synechocystis*.

14. A method according to Claim 11 wherein filamentous fungi are selected from the group consisting of *Aspergillus* and *Arthrotrys*.

30 15. A method for quantitating the amount of protein specifying RNA contained within a genome comprising:

- (a) providing a comprehensive micro-array comprising a multiplicity of open reading frames synthesized from genomic DNA comprised in a prokaryotic or eukaryotic organism;
- 35 (b) generating a set of fluorescent cDNA from total or poly-adenylated RNA isolated from the prokaryotic or eukaryotic organism of step (a);
- (c) generating a set of fluorescent DNA from genomic DNA isolated from the prokaryotic or eukaryotic organism of step (a);

- (d) hybridizing the fluorescent cDNA of step (b) to the comprehensive micro-array of step (a), wherein hybridization results in a first fluorescent signal generated from the fluorescent cDNA for each open reading frame;
- 5 (e) hybridizing the fluorescent DNA of step (c) to the comprehensive micro-array of step (a), wherein hybridization results in a second fluorescent signal generated from the fluorescent DNA for each open reading frame; and
- 10 (f) dividing, for each open reading from , the first fluorescent signal into the second fluorescent signal to provide a quantitated measure of the amount of protein specifying RNA for each open reading frame.

16. A method for quantitating the amount of protein specifying RNA contained within a genome comprising:

- 15 (a) providing a comprehensive micro-array comprising a multiplicity of genes synthesized from genomic DNA comprised in a prokaryotic or eukaryotic organism;
- (b) generating a set of fluorescent cDNA from total or poly-adenylated RNA isolated from the prokaryotic or eukaryotic organism of step (a);
- 20 (c) generating a set of fluorescent DNA from genomic DNA isolated from the prokaryotic or eukaryotic organism of step (a);
- (d) hybridizing the fluorescent cDNA of step (b) to the comprehensive micro-array of step (a), wherein hybridization results in a first fluorescent signal generated from the fluorescent cDNA for each gene;
- 25 (e) hybridizing the fluorescent DNA of step (c) to the comprehensive micro-array of step (a), wherein hybridization results in a second fluorescent signal generated from the fluorescent DNA for each gene; and
- 30 (f) dividing, for each open reading from , the first fluorescent signal into the second fluorescent signal to provide a quantitated measure of the amount of protein specifying RNA for each gene.

17. A method for identifying gene expression changes within a bacterial species according to either Claim 1 or 2 providing for quantitating the amount of protein specifying RNA contained within a genome according to either Claim 15 or 16.

18. A method for identifying gene expression changes within a genome according to Claim 8 providing for quantitating the amount of protein specifying RNA contained within a genome according to Claim 15 or 16.

1/5

FIG. 1A





FIG. 1B



2/5

FIG. 1C






FIG. 1D




3/5

FIG. 1E




4/5



EXPRESSION RANGE (%)
FIG. 2

5/5



Number of Slurred Genes

FIG. 3

Percentage of Expression